

PRODUCT MONOGRAPH  
INCLUDING PATIENT MEDICATION INFORMATION

**PrXGEVA<sup>®</sup>**

denosumab injection

120 mg/1.7 mL solution for injection  
Single-use Vial

Professed Standard

RANK Ligand Inhibitor  
(Bone Metabolism Regulator)

Amgen Canada Inc.  
6775 Financial Drive, Suite 100  
Mississauga, Ontario  
L5N 0A4

Date of Initial Authorization:

May 10, 2011

Date of Revision:

Dec 21, 2020

Submission Control Number: 235073

© 2011-2020 Amgen Canada Inc. All rights reserved.

## RECENT MAJOR LABEL CHANGES

Not Applicable	
----------------	--

## TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

<b>PART I: HEALTH PROFESSIONAL INFORMATION</b> .....	<b>4</b>
<b>1 INDICATIONS</b> .....	<b>4</b>
1.1 Pediatrics.....	4
1.2 Geriatrics (≥ 65 years of age).....	4
<b>2 CONTRAINDICATIONS</b> .....	<b>4</b>
<b>3 SERIOUS WARNINGS AND PRECAUTIONS BOX</b> .....	<b>4</b>
<b>4 DOSAGE AND ADMINISTRATION</b> .....	<b>5</b>
4.1 Dosing Considerations .....	5
4.2 Recommended Dose and Dosage Adjustment .....	5
4.3 Reconstitution.....	5
4.4 Administration.....	5
4.5 Missed Dose.....	5
<b>5 OVERDOSAGE</b> .....	<b>6</b>
<b>6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING</b> .....	<b>6</b>
<b>7 WARNINGS AND PRECAUTIONS</b> .....	<b>6</b>
7.1 Special Populations .....	9
7.1.1 Pregnant Women.....	9
7.1.2 Breast-feeding .....	10
7.1.3 Pediatrics.....	10
7.1.4 Geriatrics (≥ 65 years of age).....	11
<b>8 ADVERSE REACTIONS</b> .....	<b>11</b>
8.1 Adverse Reaction Overview .....	11
8.2 Clinical Trial Adverse Reactions.....	12
8.3 Less Common Clinical Trial Adverse Reactions .....	33
8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data .....	39
8.5 Post-Market Adverse Reactions.....	39
<b>9 DRUG INTERACTIONS</b> .....	<b>39</b>
9.1 Serious Drug Interactions .....	39
9.2 Drug Interactions Overview .....	39

9.3	Drug-Behavioural Interactions .....	39
9.4	Drug-Drug Interactions .....	39
9.5	Drug-Food Interactions.....	40
9.6	Drug-Herb Interactions .....	40
9.7	Drug-Laboratory Test Interactions.....	40
<b>10</b>	<b>CLINICAL PHARMACOLOGY .....</b>	<b>40</b>
10.1	Mechanism of Action .....	40
10.2	Pharmacodynamics .....	42
10.3	Pharmacokinetics .....	42
<b>11</b>	<b>STORAGE, STABILITY AND DISPOSAL.....</b>	<b>44</b>
<b>12</b>	<b>SPECIAL HANDLING INSTRUCTIONS.....</b>	<b>44</b>
	<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>45</b>
<b>13</b>	<b>PHARMACEUTICAL INFORMATION.....</b>	<b>45</b>
<b>14</b>	<b>CLINICAL TRIALS .....</b>	<b>45</b>
14.1	Trial Design and Study Demographics .....	45
14.2	Study Results .....	48
14.3	Comparative Bioavailability Studies .....	52
14.4	Immunogenicity .....	52
<b>15</b>	<b>MICROBIOLOGY.....</b>	<b>52</b>
<b>16</b>	<b>NON-CLINICAL TOXICOLOGY .....</b>	<b>52</b>
	<b>PATIENT MEDICATION INFORMATION .....</b>	<b>60</b>

## PART I: HEALTH PROFESSIONAL INFORMATION

### 1 INDICATIONS

XGEVA (denosumab injection) is indicated for:

- reducing the risk of developing skeletal-related events in patients with multiple myeloma and in patients with bone metastases from breast cancer, prostate cancer, non-small cell lung cancer, and other solid tumours.
- treatment of adults and skeletally mature adolescents with giant cell tumour of bone that is unresectable or where surgical resection is likely to result in severe morbidity (see **14 CLINICAL TRIALS**).
- treatment of hypercalcemia of malignancy that is refractory to intravenous bisphosphonate (see **14 CLINICAL TRIALS**).

#### 1.1 Pediatrics

The safety and efficacy of XGEVA have not been studied in pediatric populations other than skeletally mature adolescents (aged 13-17 years) with giant cell tumour of bone (GCTB). XGEVA is not indicated for use in pediatric patients other than skeletally mature adolescents with GCTB (see **7 WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics**).

#### 1.2 Geriatrics (≥ 65 years of age)

Of the total number of patients in the pivotal clinical studies in patients with advanced cancer, 1271 patients (44.4%) treated with XGEVA, were ≥ 65 years old. Of the total number of patients in the pivotal clinical study in patients with multiple myeloma, 387 patients (45.1%) treated with XGEVA, were > 65 years old. No overall differences in safety or efficacy were observed between older and younger patients.

### 2 CONTRAINDICATIONS

- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see **6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING**. Anaphylactic reactions have been reported (see **7 WARNINGS AND PRECAUTIONS, Hypersensitivity and 8 ADVERSE REACTIONS, Post-Market Adverse Reactions**).
- Pre-existing hypocalcemia must be corrected prior to initiating therapy with XGEVA (see **7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypocalcemia**).

### 3 SERIOUS WARNINGS AND PRECAUTIONS BOX

#### Serious Warnings and Precautions

Osteonecrosis of the jaw (ONJ) (see **7 WARNINGS AND PRECAUTIONS, Other, and 8 ADVERSE REACTIONS**)

## 4 DOSAGE AND ADMINISTRATION

### 4.1 Dosing Considerations

All patients, except those with hypercalcemia, should receive at least 500 mg calcium daily and at least 400 IU vitamin D daily.

### 4.2 Recommended Dose and Dosage Adjustment

#### ***Multiple Myeloma and Bone Metastasis from Solid Tumours***

The recommended dose of XGEVA is 120 mg administered as a single subcutaneous injection once every 4 weeks.

#### ***Giant Cell Tumour of Bone***

The recommended dose of XGEVA is 120 mg administered as a subcutaneous injection once every 4 weeks with a loading dose of 120 mg on days 8 and 15 of the first month of therapy.

#### ***Hypercalcemia of Malignancy Refractory to Intravenous Bisphosphonate***

The recommended dose of XGEVA is 120 mg administered as a subcutaneous injection once every 4 weeks with a loading dose of 120 mg on days 8 and 15 of the first month of therapy.

### 4.3 Reconstitution

Not applicable

### 4.4 Administration

Prior to administration, XGEVA may be removed from the refrigerator and brought to room temperature (up to 25°C) by standing in the original container. This generally takes 15 to 30 minutes. Do not warm XGEVA in any other way (see **11 STORAGE, STABILITY AND DISPOSAL**).

Visually inspect XGEVA for particulate matter and discolouration prior to administration. XGEVA is a clear, colourless to slightly yellow solution that may contain trace amounts of translucent to white proteinaceous particles. Do not use if the solution is discoloured or cloudy or if the solution contains many particles or foreign particulate matter.

Use a 27-gauge needle to withdraw and inject the entire contents of the vial. The vial is filled to ensure a deliverable dose of 120 mg. Do not re-enter the vial. Discard vial and any liquid remaining in the vial.

XGEVA is intended for use under the guidance and supervision of physicians who have fully familiarized themselves with the efficacy/safety profile of XGEVA. After an initial training in proper subcutaneous injection technique, patients may self-inject XGEVA if a physician determines that is appropriate and with medical follow-up as necessary.

XGEVA is intended for subcutaneous route only and should not be administered intravenously, intramuscularly, or intradermally. Administer XGEVA via subcutaneous injection in the upper arm, the upper thigh, or the abdomen.

### 4.5 Missed Dose

If a dose of XGEVA is missed, administer the injection as soon as the patient is available. Thereafter, injections should be scheduled every 4 weeks from the date of the last injection.

## 5 OVERDOSAGE

There is no experience with overdosage of XGEVA.

For management of a suspected drug overdose, contact your regional poison control centre.

## 6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

To help ensure the traceability of biologic products, health professionals should recognize the importance of recording both the brand name and the non-proprietary (active ingredient) name as well as other product-specific identifiers such as the Drug Identification Number (DIN) and the batch/lot number of the product supplied.

### Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Subcutaneous	120 mg denosumab in 1.7 mL solution in a single-use vial	Acetate, polysorbate 20, sodium hydroxide, sorbitol, and water for injection (USP).

XGEVA is a sterile, preservative-free, clear, colourless to slightly yellow solution, formulated at pH 5.2.

XGEVA is supplied in a single-use vial containing 120 mg denosumab, 4.6% sorbitol, 18 mM acetate, 0.01% polysorbate 20, water for injection (USP), and sodium hydroxide to a pH of 5.2.

XGEVA is supplied in a carton containing 1 vial.

## 7 WARNINGS AND PRECAUTIONS

Please see **3 SERIOUS WARNINGS AND PRECAUTIONS BOX**.

### General

XGEVA (denosumab) contains the same active ingredient as found in PROLIA®. Patients being treated with XGEVA should not be treated concomitantly with PROLIA.

Patients being treated with XGEVA should not be treated concomitantly with bisphosphonates.

### Carcinogenesis and Mutagenesis

Please see **16 NON-CLINICAL TOXICOLOGY**.

### Dermatologic

#### Skin Infections

An imbalance of skin infections leading to hospitalization was reported in a single placebo-controlled study of postmenopausal women with osteoporosis treated with denosumab 60 mg every 6 months (PROLIA 0.4%, placebo < 0.1%). These cases were predominantly cellulitis. In clinical trials in patients with advanced cancer treated with XGEVA or zoledronic acid, skin infections leading to hospitalization were reported more frequently in the XGEVA group (0.9%)

compared with the zoledronic acid group (0.7%). Patients should be advised to seek prompt medical attention if they develop signs or symptoms of cellulitis.

## Endocrine and Metabolism

### Hypocalcemia

XGEVA can cause severe symptomatic hypocalcemia and fatal cases have been reported. Signs and symptoms of severe hypocalcemia may include, for example, altered mental status, tetany, seizures and QTc prolongation. Pre-existing hypocalcemia must be corrected prior to initiating therapy with XGEVA. Monitor calcium levels, (i) prior to the initial dose of XGEVA, (ii) within two weeks after the initial dose, and (iii) if suspected symptoms of hypocalcemia occur. Administer adequate calcium and vitamin D, and magnesium, as necessary. Additional monitoring should be considered during therapy in patients with risk factors for hypocalcemia, or if otherwise indicated based on the clinical condition of the patient. Monitor levels more frequently when XGEVA is administered with other drugs that can also lower calcium levels. If hypocalcemia occurs while receiving XGEVA, additional short-term calcium supplementation and additional monitoring may be necessary (see **4 DOSAGE AND ADMINISTRATION**, and **8 ADVERSE REACTIONS, Hypocalcemia** and **Post-Market Adverse Reactions**).

An increased risk of hypocalcemia has been observed in patients with increasing degree of renal impairment. Patients with severe renal impairment (creatinine clearance less than 30 mL/min and/or receiving dialysis), are at a greater risk of developing hypocalcemia which is accompanied by an elevation in parathyroid hormone levels. Absence of calcium supplementation also plays a role in the increased risk of hypocalcemia development in patients with renal impairment. More frequent monitoring of calcium levels in these patients, within two weeks after administration of XGEVA, is especially important. If hypocalcemia occurs, administer adequate calcium, vitamin D, and magnesium, as necessary (see **7 WARNINGS AND PRECAUTIONS, Renal** and **8 ADVERSE REACTIONS**).

No dose adjustment is necessary in patients with renal impairment and there is no additional need for more frequent renal monitoring due to XGEVA administration (see **7 WARNINGS AND PRECAUTIONS, Renal**).

### Hypercalcemia Following Treatment Discontinuation in Patients with Giant Cell Tumour of Bone and in Patients with Growing Skeletons

Clinically significant hypercalcemia requiring hospitalization and complicated by acute renal injury has been reported in XGEVA-treated patients with GCTB and patients with growing skeletons weeks to months following treatment discontinuation. After treatment is discontinued, monitor patients for signs and symptoms of hypercalcemia, consider periodic assessment of serum calcium as clinically indicated, and reevaluate the patients' calcium and vitamin D supplementation requirements. Manage hypercalcemia as clinically appropriate (see **7 WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics** and **8 ADVERSE REACTIONS**).

### **Hepatic/Biliary/Pancreatic**

No clinical studies have been conducted to evaluate the effect of hepatic impairment on the pharmacokinetics of XGEVA.

## Hypersensitivity

Clinically significant hypersensitivity reactions including anaphylaxis have been reported with XGEVA. Symptoms have included hypotension, dyspnea, throat tightness, facial and upper airway edema, pruritus, and urticaria.

If an anaphylactic or other clinically significant allergic reaction occurs, initiate appropriate treatment immediately and discontinue further use of XGEVA (see **2 CONTRAINDICATIONS** and **8 ADVERSE REACTIONS**).

## Monitoring and Laboratory Tests

Monitor calcium levels, (i) prior to the initial dose of XGEVA, (ii) within two weeks after the initial dose, and (iii) if suspected symptoms of hypocalcemia occur. Additional monitoring should be considered during therapy in patients with risk factors for hypocalcemia, or if otherwise indicated based on the clinical condition of the patient. Calcium levels should be monitored more frequently when XGEVA is administered with other drugs that can also lower calcium levels (See **7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypocalcemia**).

## Other

### ***Osteonecrosis of the Jaw (ONJ)***

ONJ has been reported in patients treated with denosumab or bisphosphonates, another class of anti-resorptive agents. ONJ can manifest as jaw pain, osteomyelitis, osteitis, bone erosion, tooth or periodontal infection, toothache, gingival ulceration, or gingival erosion. Persistent pain or slow healing of the mouth or jaw after dental surgery may also be manifestations of ONJ.

In clinical trials, the incidence of ONJ was higher with longer duration of exposure (see **8 ADVERSE REACTIONS**). ONJ has also been diagnosed after treatment with XGEVA with the majority of cases occurring within 5 months after the last dose.

Poor oral hygiene, invasive dental procedures (eg, tooth extraction, dental implants, oral surgery), treatment with anti-angiogenic medication, local gum or oral infection were risk factors for ONJ in patients receiving XGEVA in clinical trials. Other risk factors for ONJ include infections, older age, concomitant therapies (eg, chemotherapy, corticosteroids, radiotherapy to the head and neck), smoking and previous treatment with bisphosphonates. In patients with risk factors for ONJ, an individual benefit-risk assessment should be performed before initiating therapy for XGEVA.

An oral exam should be performed by the prescriber prior to initiation of XGEVA treatment and a dental examination with appropriate preventive dentistry is recommended prior to treatment with XGEVA, especially in patients with risk factors for ONJ. Good oral hygiene practices should be maintained during treatment with XGEVA. Patients should receive routine dental check-ups, and immediately report any oral symptoms such as dental mobility, pain or swelling during treatment with XGEVA.

While on treatment, patients should avoid invasive dental procedures. For patients in whom invasive dental procedures cannot be avoided, the clinical judgment of the treating physician should guide the management plan of each patient based on individual benefit-risk assessment.

Patients who are suspected of having or who develop ONJ while on XGEVA should receive care by a dentist or an oral surgeon. In these patients, extensive dental surgery to treat ONJ may exacerbate the condition. In patients who develop ONJ during treatment with XGEVA, a



temporary interruption of treatment should be considered based on individual benefit-risk assessment until the condition resolves.

### ***Atypical Femoral Fractures***

Atypical femoral fracture has been reported with XGEVA. Atypical femoral fractures may occur with little or no trauma in the subtrochanteric and diaphyseal regions of the femur and may be bilateral. Specific radiographic findings characterize these events. Atypical femoral fractures have also been reported in patients with certain comorbid conditions (eg, vitamin D deficiency, rheumatoid arthritis, hypophosphatasia) and with use of certain pharmaceutical agents (eg, bisphosphonates, glucocorticoids, proton pump inhibitors). These events have also occurred without antiresorptive therapy. During XGEVA treatment, patients should be advised to report new or unusual thigh, hip, or groin pain. Patients presenting with such symptoms should be evaluated for an incomplete femoral fracture, and the contralateral femur should also be examined.

### ***Multiple Vertebral Fractures (MVF) Following Treatment Discontinuation***

Multiple vertebral fractures (MVF), not due to bone metastases, may occur following discontinuation of treatment with XGEVA, particularly in patients with risk factors such as osteoporosis or prior fractures.

Advise patients not to interrupt XGEVA therapy without their physician's advice. When XGEVA treatment is discontinued, evaluate the individual patient's risk for vertebral fractures.

## **Renal**

Two clinical trials were conducted in subjects without cancer and with varying degrees of renal function. In one study, subjects (N = 55) with varying degrees of renal function (ranging from normal through end-stage renal disease requiring dialysis) received a single 60 mg subcutaneous dose of denosumab. In a second study, patients (N = 32) with severe renal impairment (creatinine clearance less than 30 mL/minute and/or on dialysis) were given two 120 mg subcutaneous doses of denosumab. In both studies, there was a greater risk of developing hypocalcemia with increasing degree of renal impairment, and in the absence of, or inadequate calcium supplementation. The development of hypocalcemia in patients with severe renal impairment (creatinine clearance less than 30 mL/min and/or receiving dialysis) was accompanied by an elevation in parathyroid hormone levels.

No dose adjustment is necessary in patients with renal impairment and there is no additional need for more frequent renal monitoring due to XGEVA administration. Adequate intake of calcium and vitamin D is important in patients with severe renal impairment or receiving dialysis (see **7 WARNINGS AND PRECAUTIONS**, **8 ADVERSE REACTIONS**, and **10 CLINICAL PHARMACOLOGY**).

## **7.1 Special Populations**

### **7.1.1 Pregnant Women**

The safety and efficacy of XGEVA in pregnant women have not been established.

Denosumab is not recommended for use in pregnant women.

At area under the curve (AUC) exposures up to 16-fold higher than the human exposure (120 mg every 4 weeks), denosumab showed no evidence of impaired fertility in female cynomolgus monkeys.

In a study of cynomolgus monkeys dosed with denosumab during the period equivalent to the first trimester at AUC exposures up to 10-fold higher than the human dose (120 mg every 4 weeks), there was no evidence of maternal or fetal harm. In this study, fetal lymph nodes were not examined.

In another study, in utero denosumab exposure in cynomolgus monkeys at 50 mg/kg body weight every 4 weeks, from gestation day 20 through to parturition resulted in increased fetal loss, stillbirths and postnatal mortality. Findings in the infants included skeletal abnormalities resulting from impaired bone resorption during rapid growth, reduced bone strength and treatment-related bone fractures; reduced hematopoiesis; tooth malalignment and dental dysplasia (in the absence of adverse effects on tooth eruption); absence of peripheral lymph nodes; and decreased neonatal growth. There was no evidence of maternal toxicity. Maternal mammary gland development was normal.

In genetically engineered mice in which the gene for RANK ligand (RANKL) has been deleted (a “knockout mouse”), the absence of RANKL caused fetal lymph node agenesis and led to postnatal impairment of dentition and bone growth. Pregnant RANKL knockout mice also showed altered maturation of the maternal mammary gland, leading to impaired lactation postpartum (see **16 NON-CLINICAL TOXICOLOGY**).

Women should be advised not to become pregnant during XGEVA therapy. Advise females of reproductive potential to use highly effective contraception during therapy, and for at least 5 months after the last dose of XGEVA.

### **7.1.2 Breast-feeding**

It is not known whether XGEVA is excreted into human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from XGEVA, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Maternal exposure to XGEVA during pregnancy may impair mammary gland development and lactation based on animal studies in pregnant mice lacking the RANK/RANKL signaling pathway which showed altered maturation of the maternal mammary gland, leading to impaired lactation postpartum (see **16 NON-CLINICAL TOXICOLOGY**).

### **7.1.3 Pediatrics**

The safety and efficacy of XGEVA have not been established in pediatric patients other than skeletally mature adolescents (aged 13-17 years) with GCTB.

XGEVA is not recommended for use in pediatric patients other than skeletally mature adolescents with GCTB. Clinically significant hypercalcemia after treatment discontinuation has been reported in the post-marketing setting in pediatric patients with growing skeletons who received denosumab for GCTB or for other unauthorized uses (see **7 WARNINGS AND PRECAUTIONS**).

XGEVA was studied in a phase 2 open-label trial that enrolled a subset of adolescent patients (10 at interim analysis and 28 at final analysis; aged 13-17 years) with GCTB who had reached skeletal maturity defined by at least 1 mature long bone (eg, closed epiphyseal growth plate of the humerus) and body weight  $\geq$  45 kg (see **1 INDICATIONS** and **14 CLINICAL TRIALS**). The adverse reaction profile appeared similar in skeletally mature adolescents and adults.

Treatment with XGEVA may impair bone growth in children with open growth plates and may inhibit eruption of dentition. In neonatal rats, inhibition of RANKL (target of XGEVA therapy)

with a construct of osteoprotegerin bound to Fc (OPG-Fc) at doses  $\leq$  10 mg/kg was associated with inhibition of bone growth and tooth eruption. Adolescent monkeys dosed with denosumab at 15 times (50 mg/kg dose) and 2.8 times (10 mg/kg dose) the area under the curve (AUC) exposure in adult humans dosed at 120 mg subcutaneously every 4 weeks had abnormal growth plates, considered to be consistent with the pharmacological activity of denosumab. In neonatal cynomolgus monkeys exposed in utero to denosumab at 50 mg/kg, there was increased postnatal mortality; skeletal abnormalities resulting from impaired bone resorption during rapid growth, reduced bone strength and treatment-related bone fractures; reduced hematopoiesis; tooth malalignment and dental dysplasia (in the absence of adverse effects on tooth eruption); absence of peripheral lymph nodes; and decreased neonatal growth. Following a recovery period from birth out to 6 months of age, findings still observed were mildly reduced bone length (femoral, vertebral, jaw), reduced cortical thickness with associated reduced strength; extramedullary hematopoiesis; dental dysplasia; and the absence or decreased size of some lymph nodes. One infant had minimal to moderate mineralization in multiple tissues (see **16 NON-CLINICAL TOXICOLOGY**).

#### 7.1.4 Geriatrics ( $\geq$ 65 years of age)

Of the total number of patients in the pivotal clinical studies in patients with advanced cancer, 1271 patients (44.4%) treated with XGEVA were  $\geq$  65 years old. Of the total number of patients in the pivotal clinical study in patients with multiple myeloma, 387 patients (45.1%) treated with XGEVA, were  $>$  65 years old. No overall differences in safety or efficacy were observed between older and younger patients.

## 8 ADVERSE REACTIONS

### 8.1 Adverse Reaction Overview

The following adverse reactions are discussed below and elsewhere in the Product Monograph:

- Hypocalcemia (see **7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypocalcemia**)
- Osteonecrosis of the jaw (see **7 WARNINGS AND PRECAUTIONS, Other, Osteonecrosis of the Jaw (ONJ)**)
- Multiple vertebral fractures following treatment discontinuation (see **7 WARNINGS AND PRECAUTIONS, Other, Multiple Vertebral Fractures (MVF) Following Treatment Discontinuation**)

The most common adverse reactions in patients with bone metastasis from solid tumours receiving XGEVA (per-patient incidence  $\geq$  25%) were fatigue/asthenia, hypophosphatemia, and nausea.

The most common serious adverse reaction in patients with bone metastasis from solid tumours receiving XGEVA was dyspnea.

The most common adverse reactions in patients with bone metastasis from solid tumours resulting in discontinuation of XGEVA were osteonecrosis and hypocalcemia.

The most common adverse reactions in patients with multiple myeloma (per-patient incidence  $\geq$  10%) were back pain, hypocalcemia, arthralgia, and bone pain.

The most common serious adverse reactions in patients with multiple myeloma (per-patient incidence  $\geq$  1%) were back pain and hypocalcemia.

The most common adverse reactions resulting in discontinuation of XGEVA in patients with multiple myeloma (per-patient incidence  $\geq 1\%$ ) was osteonecrosis of the jaw.

In the initial integrated analysis (final analysis of Study 5 [20040215] and interim analysis of Study 6 [20062004]), the most common adverse reactions in patients with GCTB receiving XGEVA (per-patient incidence  $\geq 10\%$ ) were arthralgia, headache, nausea, fatigue, back pain, and pain in extremity. In the final integrated analysis (final analyses of both Studies 5 and 6), the following additional common adverse reactions were observed (per-patient incidence  $\geq 10\%$ ): nasopharyngitis, musculoskeletal pain, toothache, vomiting, hypophosphatemia, constipation, diarrhea, and cough.

In the initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6), the most common serious adverse reactions in patients with GCTB receiving XGEVA (per-patient incidence of 0.7%) were osteonecrosis of the jaw and osteomyelitis. In the final integrated analysis (final analyses of both Studies 5 and 6), the following most common serious adverse reactions were observed (per patient incidence of  $\geq 0.9\%$ ): osteonecrosis of the jaw, bone giant cell tumour, anemia, pneumonia and back pain.

In the initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6), the most common adverse reactions in patients with GCTB receiving XGEVA resulting in discontinuation of XGEVA (per-patient incidence of 0.7%) were osteonecrosis of the jaw and tooth abscess or tooth infection. In the final integrated analysis (final analyses of both Studies 5 and 6), the most common adverse reaction in patients with GCTB receiving XGEVA resulting in discontinuation of XGEVA (per-patient incidence  $\geq 0.9\%$ ) was osteonecrosis of the jaw.

## 8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse drug reactions in real-world use.

An **adverse reaction** is an unintended event, reasonably associated with the use of a drug. The data presented below, however, reflect **adverse events**, which do not necessarily have a causal relationship to the drug.

### Bone Metastasis from Solid Tumours

The safety of XGEVA was evaluated in three randomized, double-blind, double-dummy trials (see **14 CLINICAL TRIALS**) in which a total of 2841 patients with bone metastasis from prostate cancer, breast cancer, or other solid tumours, or lytic bony lesions from multiple myeloma received at least one dose of XGEVA. In Studies 1, 2, and 3, patients were randomized to receive either 120 mg of XGEVA every 4 weeks as a subcutaneous injection or 4 mg (dose adjusted for reduced renal function) of zoledronic acid every 4 weeks by intravenous (IV) infusion. Entry criteria included serum calcium (corrected) from 8 to 11.5 mg/dL (2 to 2.9 mmol/L) and creatinine clearance 30 mL/min or greater. Patients who had received IV bisphosphonates were excluded, as were patients with prior history of ONJ or osteomyelitis of the jaw, an active dental or jaw condition requiring oral surgery, non-healed dental/oral surgery, or any planned invasive dental procedure. During the study, serum chemistries including calcium and phosphorus were monitored every 4 weeks. Calcium and vitamin D supplementation was recommended but not required.

The median duration of exposure to XGEVA was 12 months (range: 0.1 – 41) and median duration on-study was 13 months (range: 0.1 – 41). Of patients who received XGEVA, 46%

were female. Eighty-five percent were White, 5% Hispanic/Latino, 6% Asian, and 3% Black. The median age was 63 years (range: 18 – 93). Seventy-five percent of patients who received XGEVA received concomitant chemotherapy.

The adverse events occurring during the studies were generally of a type and frequency expected in patients with cancer and bone metastases, many of whom were undergoing antineoplastic therapy. Table 1 describes adverse events occurring in  $\geq 1\%$  of patients in these studies.

**Table 1. Percentage of Patients with Adverse Events  $\geq 1\%$  Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
<b>BLOOD AND LYMPHATIC SYSTEM DISORDERS</b>		
Anemia	771 (27.1)	859 (30.3)
Neutropenia	277 (9.8)	278 (9.8)
Thrombocytopenia	216 (7.6)	199 (7.0)
Leukopenia	165 (5.8)	177 (6.2)
Febrile neutropenia	58 (2.0)	72 (2.5)
Pancytopenia	29 (1.0)	34 (1.2)
<b>CARDIAC DISORDERS</b>		
Tachycardia	79 (2.8)	74 (2.6)
Cardiac failure	49 (1.7)	51 (1.8)
Atrial fibrillation	43 (1.5)	38 (1.3)
Palpitations	30 (1.1)	25 (0.9)
<b>EAR AND LABYRINTH DISORDERS</b>		
Vertigo	62 (2.2)	85 (3.0)
<b>EYE DISORDERS</b>		
Lacrimation increased	59 (2.1)	46 (1.6)
Vision blurred	53 (1.9)	48 (1.7)
Conjunctivitis	31 (1.1)	37 (1.3)
<b>GASTROINTESTINAL DISORDERS</b>		
Nausea	876 (30.8)	895 (31.6)
Constipation	603 (21.2)	670 (23.6)
Diarrhea	577 (20.3)	530 (18.7)
Vomiting	566 (19.9)	570 (20.1)
Abdominal pain	292 (10.3)	280 (9.9)
Abdominal pain upper	167 (5.9)	164 (5.8)
Stomatitis	146 (5.1)	115 (4.1)
Dyspepsia	132 (4.6)	147 (5.2)
Toothache	108 (3.8)	80 (2.8)
Ascites	68 (2.4)	53 (1.9)
Dysphagia	66 (2.3)	63 (2.2)
Abdominal distension	56 (2.0)	47 (1.7)
Dry mouth	53 (1.9)	57 (2.0)

**Table 1. Percentage of Patients with Adverse Events  $\geq$  1% Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
Gastritis	51 (1.8)	59 (2.1)
Hemorrhoids	50 (1.8)	52 (1.8)
Gastroesophageal reflux disease	49 (1.7)	50 (1.8)
Flatulence	42 (1.5)	38 (1.3)
Gingival pain	36 (1.3)	29 (1.0)
Rectal hemorrhage	32 (1.1)	37 (1.3)
Abdominal discomfort	30 (1.1)	26 (0.9)
Gingivitis	30 (1.1)	26 (0.9)
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>		
Fatigue	769 (27.1)	766 (27.0)
Asthenia	607 (21.4)	621 (21.9)
Edema peripheral	472 (16.6)	462 (16.3)
Pyrexia	409 (14.4)	562 (19.8)
Chest pain	263 (9.3)	247 (8.7)
Pain	222 (7.8)	243 (8.6)
General physical health deterioration	131 (4.6)	135 (4.8)
Mucosal inflammation	123 (4.3)	133 (4.7)
Edema	71 (2.5)	100 (3.5)
Chills	55 (1.9)	115 (4.1)
Influenza like illness	43 (1.5)	83 (2.9)
Malaise	41 (1.4)	36 (1.3)
Multi-organ failure	37 (1.3)	35 (1.2)
Gait disturbance	33 (1.2)	35 (1.2)
Face edema	29 (1.0)	17 (0.6)
Chest discomfort	26 (0.9)	31 (1.1)
Disease progression	25 (0.9)	31 (1.1)
<b>HEPATOBIILIARY DISORDERS</b>		
Hepatic failure	41 (1.4)	31 (1.1)
Hepatic function abnormal	37 (1.3)	28 (1.0)
Jaundice	29 (1.0)	21 (0.7)
<b>INFECTIONS AND INFESTATIONS</b>		
Urinary tract infection	220 (7.7)	262 (9.2)
Nasopharyngitis	149 (5.2)	163 (5.7)
Pneumonia	147 (5.2)	130 (4.6)
Bronchitis	124 (4.4)	103 (3.6)
Influenza	118 (4.2)	97 (3.4)
Upper respiratory tract infection	110 (3.9)	116 (4.1)
Oral candidiasis	81 (2.9)	74 (2.6)
Sinusitis	70 (2.5)	50 (1.8)
Herpes zoster	54 (1.9)	49 (1.7)

**Table 1. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
Cellulitis	51 (1.8)	47 (1.7)
Rhinitis	46 (1.6)	40 (1.4)
Cystitis	44 (1.5)	48 (1.7)
Respiratory tract infection	40 (1.4)	36 (1.3)
Sepsis	37 (1.3)	34 (1.2)
Pharyngitis	35 (1.2)	41 (1.4)
Gastroenteritis	30 (1.1)	26 (0.9)
Oral herpes	30 (1.1)	24 (0.8)
Tooth abscess	29 (1.0)	16 (0.6)
Lower respiratory tract infection	20 (0.7)	29 (1.0)
<b>INJURY, POISONING AND PROCEDURAL COMPLICATIONS</b>		
Rib fracture	158 (5.6)	166 (5.9)
Thoracic vertebral fracture	149 (5.2)	154 (5.4)
Lumbar vertebral fracture	107 (3.8)	111 (3.9)
Contusion	61 (2.1)	57 (2.0)
Fall	54 (1.9)	48 (1.7)
Femur fracture	33 (1.2)	37 (1.3)
<b>INVESTIGATIONS</b>		
Hypophosphatemia (laboratory-derived)	912 (32.1)	555 (19.6)
Weight decreased	330 (11.6)	332 (11.7)
Blood creatinine increased	105 (3.7)	134 (4.7)
Aspartate aminotransferase increased	76 (2.7)	95 (3.3)
Blood alkaline phosphatase increased	74 (2.6)	76 (2.7)
Alanine aminotransferase increased	62 (2.2)	82 (2.9)
Hemoglobin decreased	56 (2.0)	60 (2.1)
Weight increased	48 (1.7)	55 (1.9)
Platelet count decreased	39 (1.4)	36 (1.3)
Prostatic specific antigen increased	37 (1.3)	19 (0.7)
<b>METABOLISM AND NUTRITION DISORDERS</b>		
Decreased appetite	656 (23.1)	694 (24.5)
Hypocalcemia	265 (9.3)	134 (4.7)
Dehydration	179 (6.3)	164 (5.8)
Hypokalemia	130 (4.6)	156 (5.5)
Hyperglycemia	108 (3.8)	107 (3.8)
Hypophosphatemia	61 (2.1)	32 (1.1)
Hypomagnesemia	56 (2.0)	46 (1.6)
Hyponatremia	50 (1.8)	64 (2.3)
Hypoalbuminemia	48 (1.7)	44 (1.6)
Hyperkalemia	45 (1.6)	50 (1.8)
Hypercalcemia	39 (1.4)	51 (1.8)

**Table 1. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
Cachexia	35 (1.2)	37 (1.3)
Hypoglycemia	29 (1.0)	32 (1.1)
<b>MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS</b>		
Back pain	718 (25.3)	747 (26.3)
Arthralgia	570 (20.1)	632 (22.3)
Bone pain	564 (19.9)	639 (22.5)
Pain in extremity	524 (18.4)	550 (19.4)
Musculoskeletal pain	357 (12.6)	385 (13.6)
Musculoskeletal chest pain	186 (6.5)	188 (6.6)
Myalgia	150 (5.3)	195 (6.9)
Neck pain	125 (4.4)	144 (5.1)
Muscle spasms	121 (4.3)	96 (3.4)
Muscular weakness	111 (3.9)	140 (4.9)
Pain in jaw	108 (3.8)	83 (2.9)
Osteonecrosis	52 (1.8)	34 (1.2)
Groin pain	48 (1.7)	63 (2.2)
Flank pain	31 (1.1)	34 (1.2)
Musculoskeletal stiffness	30 (1.1)	45 (1.6)
Joint swelling	29 (1.0)	25 (0.9)
<b>NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS)</b>		
Metastases to central nervous system	137 (4.8)	122 (4.3)
Malignant neoplasm progression	130 (4.6)	129 (4.5)
Metastases to liver	103 (3.6)	88 (3.1)
Metastases to bone	94 (3.3)	97 (3.4)
Prostate cancer	47 (1.7)	66 (2.3)
Metastases to spine	40 (1.4)	41 (1.4)
Metastases to lung	33 (1.2)	32 (1.1)
<b>NERVOUS SYSTEM DISORDERS</b>		
Headache	360 (12.7)	382 (13.5)
Dizziness	232 (8.2)	254 (9.0)
Paresthesia	168 (5.9)	204 (7.2)
Neuropathy peripheral	147 (5.2)	142 (5.0)
Hypoesthesia	109 (3.8)	118 (4.2)
Dysgeusia	104 (3.7)	102 (3.6)
Spinal cord compression	96 (3.4)	118 (4.2)
Peripheral sensory neuropathy	89 (3.1)	86 (3.0)
Somnolence	57 (2.0)	69 (2.4)
Syncope	50 (1.8)	50 (1.8)
Lethargy	44 (1.5)	51 (1.8)
Sciatica	37 (1.3)	40 (1.4)



**Table 1. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
Tremor	27 (1.0)	46 (1.6)
Convulsion	27 (1.0)	32 (1.1)
Neuralgia	27 (1.0)	30 (1.1)
<b>PSYCHIATRIC DISORDERS</b>		
Insomnia	302 (10.6)	324 (11.4)
Anxiety	196 (6.9)	184 (6.5)
Depression	186 (6.5)	182 (6.4)
Confusional state	87 (3.1)	87 (3.1)
Agitation	20 (0.7)	35 (1.2)
<b>RENAL AND URINARY DISORDERS</b>		
Hematuria	115 (4.0)	118 (4.2)
Urinary retention	112 (3.9)	109 (3.8)
Dysuria	111 (3.9)	102 (3.6)
Renal failure	74 (2.6)	104 (3.7)
Pollakiuria	59 (2.1)	69 (2.4)
Hydronephrosis	56 (2.0)	47 (1.7)
Urinary incontinence	40 (1.4)	54 (1.9)
Renal failure acute	34 (1.2)	44 (1.6)
Renal impairment	26 (0.9)	34 (1.2)
Nocturia	23 (0.8)	36 (1.3)
<b>REPRODUCTIVE SYSTEM AND BREAST DISORDERS</b>		
Pelvic pain	80 (2.8)	79 (2.8)
Breast pain	28 (1.0)	31 (1.1)
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>		
Dyspnea	585 (20.6)	507 (17.9)
Cough	437 (15.4)	419 (14.8)
Pleural effusion	153 (5.4)	137 (4.8)
Epistaxis	109 (3.8)	107 (3.8)
Oropharyngeal pain	96 (3.4)	81 (2.9)
Respiratory failure	96 (3.4)	78 (2.8)
Dyspnea exertional	58 (2.0)	53 (1.9)
Pulmonary embolism	57 (2.0)	65 (2.3)
Hemoptysis	47 (1.7)	51 (1.8)
Dysphonia	46 (1.6)	49 (1.7)
Productive cough	35 (1.2)	37 (1.3)
Nasal congestion	30 (1.1)	23 (0.8)
Hypoxia	29 (1.0)	21 (0.7)
Rhinorrhea	24 (0.8)	31 (1.1)

**Table 1. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Advanced Malignancies Involving Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N = 2841) n (%)</b>	<b>Zoledronic Acid (N = 2836) n (%)</b>
<b>SKIN AND SUBCUTANEOUS TISSUE DISORDERS</b>		
Alopecia	265 (9.3)	266 (9.4)
Rash	193 (6.8)	201 (7.1)
Pruritus	107 (3.8)	111 (3.9)
Palmar-plantar erythrodysesthesia syndrome	101 (3.6)	109 (3.8)
Nail disorder	66 (2.3)	72 (2.5)
Hyperhidrosis	66 (2.3)	36 (1.3)
Erythema	64 (2.3)	70 (2.5)
Dry skin	58 (2.0)	60 (2.1)
Night sweats	32 (1.1)	33 (1.2)
Dermatitis	31 (1.1)	20 (0.7)
Skin ulcer	30 (1.1)	19 (0.7)
<b>SURGICAL AND MEDICAL PROCEDURES</b>		
Tooth extraction	43 (1.5)	24 (0.8)
<b>VASCULAR DISORDERS</b>		
Hypertension	148 (5.2)	153 (5.4)
Hypotension	112 (3.9)	99 (3.5)
Hot flush	95 (3.3)	98 (3.5)
Deep vein thrombosis	51 (1.8)	55 (1.9)
Lymphedema	47 (1.7)	43 (1.5)
Phlebitis	31 (1.1)	31 (1.1)

N = Number of subjects who received ≥ 1 active dose of investigational product

n = Number of subjects reporting ≥ 1 event

Includes only treatment-emergent adverse events

System organ classes are sorted alphabetically and preferred terms are sorted by descending order of frequency in the XGEVA group and coded using MedDRA version 12.1

## Multiple Myeloma

In Study 4, XGEVA was evaluated in an international, randomized (1:1), double blind, active controlled study comparing XGEVA with zoledronic acid in patients with newly diagnosed multiple myeloma (see **14 CLINICAL TRIALS**). In this study, 1718 multiple myeloma patients with at least 1 bone lesion were randomized to receive 120 mg XGEVA subcutaneously every 4 weeks or 4 mg zoledronic acid intravenously (IV) every 4 weeks (dose adjusted for renal impairment and patients with creatinine clearance less than 30 mL/min were excluded based on zoledronic acid prescribing information).

The number of patients across both study arms with ISS stage I, stage II, and stage III at diagnosis were 32.4%, 38.2%, and 29.3%, respectively. In the XGEVA group, 46% were female, 82% were White, 12% Asian, and 4% Black. The median age was 63 years (range: 29 - 91). The median duration of exposure to XGEVA was 16 months (range: 1 – 50) and median duration on study was 17 months (range: 0 – 49). The median number of doses administered was 16 for XGEVA and 15 for zoledronic acid.

Table 2 describes adverse events occurring in  $\geq 1\%$  of patients in this study.

**Table 2. Percentage of Patients with Adverse Events  $\geq 1\%$  Reported in Patients with Multiple Myeloma by System Organ Class\***

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA 120 mg Q4W (N = 850) n (%)</b>	<b>Zoledronic Acid 4 mg Q4W (N = 852) n (%)</b>
<b>BLOOD AND LYMPHATIC SYSTEM DISORDERS</b>		
Anemia	189 (22.2)	180 (21.1)
Neutropenia	160 (18.8)	155 (18.2)
Thrombocytopenia	160 (18.8)	137 (16.1)
Febrile neutropenia	102 (12.0)	98 (11.5)
Leukopenia	42 (4.9)	36 (4.2)
<b>CARDIAC DISORDERS</b>		
Cardiac failure	13 (1.5)	8 (0.9)
Palpitations	13 (1.5)	4 (0.5)
<b>EAR AND LABYRINTH DISORDERS</b>		
Vertigo	16 (1.9)	7 (0.8)
Ear pain	14 (1.6)	4 (0.5)
Tinnitus	11 (1.3)	9 (1.1)
<b>EYE DISORDERS</b>		
Vision blurred	30 (3.5)	19 (2.2)
Lacrimation increased	9 (1.1)	6 (0.7)
<b>GASTROINTESTINAL DISORDERS</b>		
Diarrhea	285 (33.5)	276 (32.4)
Nausea	268 (31.5)	259 (30.4)
Abdominal pain	66 (7.8)	50 (5.9)
Stomatitis	59 (6.9)	50 (5.9)
Toothache	41 (4.8)	34 (4.0)
Abdominal distension	20 (2.4)	20 (2.3)
Abdominal discomfort	18 (2.1)	11 (1.3)
Oral pain	16 (1.9)	13 (1.5)
Mouth ulceration	15 (1.8)	12 (1.4)
Dysphagia	12 (1.4)	10 (1.2)
Flatulence	11 (1.3)	8 (0.9)
Gastritis	9 (1.1)	9 (1.1)
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>		
Edema peripheral	148 (17.4)	135 (15.8)
Mucosal inflammation	72 (8.5)	49 (5.8)
Pain	41 (4.8)	36 (4.2)
Chest pain	29 (3.4)	25 (2.9)
Influenza like illness	28 (3.3)	26 (3.1)
Peripheral swelling	27 (3.2)	17 (2.0)
Non-cardiac chest pain	13 (1.5)	12 (1.4)

**Table 2. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Multiple Myeloma by System Organ Class\***

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA 120 mg Q4W (N = 850) n (%)</b>	<b>Zoledronic Acid 4 mg Q4W (N = 852) n (%)</b>
General physical health deterioration	11 (1.3)	9 (1.1)
Gait disturbance	10 (1.2)	8 (0.9)
Malaise	10 (1.2)	8 (0.9)
Face Edema	10 (1.2)	7 (0.8)
<b>IMMUNE SYSTEM DISORDERS</b>		
Hypersensitivity	9 (1.1)	8 (0.9)
<b>INFECTIONS AND INFESTATIONS</b>		
Upper respiratory tract infection	130 (15.3)	109 (12.8)
Pneumonia	98 (11.5)	97 (11.4)
Nasopharyngitis	72 (8.5)	62 (7.3)
Bronchitis	67 (7.9)	67 (7.9)
Urinary tract infection	59 (6.9)	52 (6.1)
Respiratory tract infection	56 (6.6)	49 (5.8)
Lower respiratory tract infection	24 (2.8)	24 (2.8)
Cellulitis	13 (1.5)	13 (1.5)
Rhinitis	13 (1.5)	11 (1.3)
Periodontitis	12 (1.4)	12 (1.4)
Gingivitis	12 (1.4)	11 (1.3)
Pharyngitis	11 (1.3)	10 (1.2)
Viral infection	11 (1.3)	9 (1.1)
Folliculitis	10 (1.2)	10 (1.2)
Hordeolum	10 (1.2)	7 (0.8)
<i>Clostridium difficile</i> colitis	10 (1.2)	6 (0.7)
Urinary tract infection bacterial	10 (1.2)	6 (0.7)
Bacterial infection	9 (1.1)	8 (0.9)
<b>INJURY, POISONING AND PROCEDURAL COMPLICATIONS</b>		
Thoracic vertebral fracture	84 (9.9)	75 (8.8)
Contusion	16 (1.9)	11 (1.3)
Cervical vertebral fracture	13 (1.5)	7 (0.8)
Pubis fracture	10 (1.2)	6 (0.7)
<b>INVESTIGATIONS</b>		
Weight decreased	59 (6.9)	56 (6.6)
Aspartate aminotransferase increased	28 (3.3)	27 (3.2)
Weight increased	28 (3.3)	22 (2.6)
White blood cell count decreased	21 (2.5)	18 (2.1)
Blood alkaline phosphatase increased	14 (1.6)	12 (1.4)
Gamma-glutamyltransferase increased	14 (1.6)	11 (1.3)
C-reactive protein increased	13 (1.5)	13 (1.5)
Blood glucose increased	9 (1.1)	6 (0.7)

**Table 2. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Multiple Myeloma by System Organ Class\***

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA 120 mg Q4W (N = 850) n (%)</b>	<b>Zoledronic Acid 4 mg Q4W (N = 852) n (%)</b>
<b>METABOLISM AND NUTRITION DISORDERS</b>		
Hypocalcemia	139 (16.4)	101 (11.9)
Hypophosphatemia	54 (6.4)	43 (5.0)
Hyperglycemia	35 (4.1)	28 (3.3)
Dehydration	33 (3.9)	28 (3.3)
Hyperkalemia	26 (3.1)	12 (1.4)
Hyponatremia	25 (2.9)	22 (2.6)
Hypoalbuminemia	23 (2.7)	15 (1.8)
Hyperuricemia	14 (1.6)	12 (1.4)
Gout	11 (1.3)	7 (0.8)
<b>MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS</b>		
Back pain	178 (20.9)	167 (19.6)
Arthralgia	115 (13.5)	112 (13.1)
Muscle spasms	53 (6.2)	41 (4.8)
Musculoskeletal chest pain	42 (4.9)	34 (4.0)
Osteonecrosis of jaw	38 (4.5)	25 (2.9)
Neck pain	27 (3.2)	23 (2.7)
Osteoarthritis	16 (1.9)	11 (1.3)
Flank pain	12 (1.4)	9 (1.1)
Joint swelling	11 (1.3)	10 (1.2)
Musculoskeletal discomfort	10 (1.2)	9 (1.1)
<b>NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS)</b>		
Plasma cell myeloma	26 (3.1)	26 (3.1)
Plasmacytoma	10 (1.2)	4 (0.5)
<b>NERVOUS SYSTEM DISORDERS</b>		
Headache	96 (11.3)	77 (9.0)
Peripheral sensory neuropathy	75 (8.8)	71 (8.3)
Hypoesthesia	47 (5.5)	38 (4.5)
Tremor	26 (3.1)	21 (2.5)
Sciatica	10 (1.2)	8 (0.9)
<b>PSYCHIATRIC DISORDERS</b>		
Insomnia	123 (14.5)	123 (14.4)
Confusional state	24 (2.8)	12 (1.4)
Delirium	9 (1.1)	4 (0.5)
<b>RENAL AND URINARY DISORDERS</b>		
Dysuria	21 (2.5)	16 (1.9)
Urinary retention	12 (1.4)	8 (0.9)
Pollakiuria	12 (1.4)	7 (0.8)

**Table 2. Percentage of Patients with Adverse Events ≥ 1% Reported in Patients with Multiple Myeloma by System Organ Class\***

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA 120 mg Q4W (N = 850) n (%)</b>	<b>Zoledronic Acid 4 mg Q4W (N = 852) n (%)</b>
Hematuria	9 (1.1)	9 (1.1)
Urinary incontinence	9 (1.1)	4 (0.5)
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>		
Oropharyngeal pain	57 (6.7)	40 (4.7)
Rhinorrhea	22 (2.6)	18 (2.1)
Pleural effusion	16 (1.9)	16 (1.9)
Pulmonary embolism	15 (1.8)	15 (1.8)
Wheezing	13 (1.5)	7 (0.8)
Dysphonia	12 (1.4)	8 (0.9)
Sinus congestion	10 (1.2)	2 (0.2)
Pneumonitis	9 (1.1)	0 (0.0)
<b>SKIN AND SUBCUTANEOUS TISSUE DISORDERS</b>		
Rash	119 (14.0)	96 (11.3)
Alopecia	54 (6.4)	48 (5.6)
Dry skin	26 (3.1)	17 (2.0)
Erythema	23 (2.7)	21 (2.5)
Night sweats	16 (1.9)	10 (1.2)
Hyperhidrosis	14 (1.6)	13 (1.5)
Rash generalized	14 (1.6)	12 (1.4)
Rash maculo-papular	14 (1.6)	10 (1.2)
Urticaria	11 (1.3)	5 (0.6)
Eczema	9 (1.1)	9 (1.1)
Skin lesion	9 (1.1)	9 (1.1)
Petechiae	9 (1.1)	5 (0.6)
Dermatitis allergic	9 (1.1)	1 (0.1)
<b>VASCULAR DISORDERS</b>		
Hypertension	61 (7.2)	60 (7.0)
Hypotension	49 (5.8)	33 (3.9)
Deep vein thrombosis	28 (3.3)	23 (2.7)
Orthostatic hypotension	12 (1.4)	11 (1.3)

\*Adverse Events Occurring in ≥ 1% Subject Incidence in the XGEVA Group and Higher Than Zoledronic Acid Group by System Organ Class and Preferred Term

N = Number of subjects who received ≥ 1 active dose of investigational product

n = Number of subjects reporting ≥ 1 event

Includes only treatment-emergent adverse events

System organ classes are sorted alphabetically and preferred terms are sorted by descending order of frequency in the XGEVA group and coded using MedDRA version 19.0

## **Giant Cell Tumour of Bone**

The safety of XGEVA was evaluated in two phase 2 open-label, single arm studies (Study 5 and 6) (see **14 CLINICAL TRIALS**) in which patients (n=304 at initial integrated analysis and n=548 at final integrated analysis) with GCTB received at least 1 dose of XGEVA. Patients received 120 mg XGEVA subcutaneously every 4 weeks with a loading dose of 120 mg on days 8 and 15. In the initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6), of the 304 patients who received XGEVA, 147 patients were treated with XGEVA for  $\geq 1$  year, 46 patients for  $\geq 2$  years, and 15 patients for  $\geq 3$  years. The median number of doses received was 14 (range: 1 to 60 doses) and the median number of months on study was 11 (range: 0 to 54 months).

Fifty-eight percent of the enrolled subjects were women. The majority of subjects were white (80.3%). The median (range) age was 33 (13 to 83) years; 10 subjects were skeletally mature adolescents (aged 13 to 17 years).

At the final integrated analysis of both Studies 5 and 6, of the 548 patients who received XGEVA, 467 patients were treated with XGEVA for  $\geq 1$  year, 323 patients for  $\geq 2$  years, and 255 patients for  $\geq 3$  years. The median number of doses received was 33 (range: 4 to 138 doses) and the median number of months on study was 60 (range: 0 to 140 months) in the final integrated analysis.

Fifty-seven percent of the enrolled subjects were women. The majority of subjects were white (81.9%). The median (range) age was 33 (13 to 83) years; 28 subjects were skeletally mature adolescents (aged 13 to 17 years).

Table 3 describes adverse events at initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6) occurring in  $\geq 1\%$  of patients in these studies.

**Table 3. Percentage of Patients with Adverse Events  $\geq 1\%$  Reported in Patients with Giant Cell Tumour of Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N=304) n (%)</b>
<b>BLOOD AND LYMPHATIC SYSTEM DISORDERS</b>	
Anemia	12 (3.9)
<b>CARDIAC DISORDERS</b>	
Palpitations	5 (1.6)
Tachycardia	4 (1.3)
<b>EAR AND LABYRINTH DISORDERS</b>	
Vertigo	10 (3.3)
<b>GASTROINTESTINAL DISORDERS</b>	
Nausea	54 (17.8)
Vomiting	28 (9.2)
Constipation	22 (7.2)
Diarrhea	21 (6.9)
Toothache	17 (5.6)
Abdominal pain	16 (5.3)
Abdominal pain upper	11 (3.6)
Dyspepsia	8 (2.6)

**Table 3. Percentage of Patients with Adverse Events  $\geq$  1% Reported in Patients with Giant Cell Tumour of Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N=304) n (%)</b>
Abdominal distension	6 (2.0)
Dental caries	5 (1.6)
Stomatitis	5 (1.6)
Tooth disorder	5 (1.6)
Abdominal discomfort	4 (1.3)
Dry mouth	4 (1.3)
Gastroesophageal reflux disease	4 (1.3)
<b>GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS</b>	
Fatigue	51 (16.8)
Edema peripheral	24 (7.9)
Non-cardiac chest pain	16 (5.3)
Pyrexia	14 (4.6)
Asthenia	12 (3.9)
Influenza like illness	11 (3.6)
Local swelling	8 (2.6)
Pain	7 (2.3)
Chills	5 (1.6)
Injection site hematoma	4 (1.3)
<b>INFECTIONS AND INFESTATIONS</b>	
Nasopharyngitis	24 (7.9)
Upper respiratory tract infection	23 (7.6)
Urinary tract infection	11 (3.6)
Influenza	9 (3.0)
Gastroenteritis	8 (2.6)
Sinusitis	7 (2.3)
Cystitis	6 (2.0)
Bronchitis	5 (1.6)
Tooth abscess	5 (1.6)
Pharyngitis	4 (1.3)
Tooth infection	4 (1.3)
<b>INJURY, POISONING AND PROCEDURAL COMPLICATIONS</b>	
Procedural pain	13 (4.3)
Contusion	4 (1.3)
<b>INVESTIGATIONS</b>	
Weight increased	19 (6.3)
Weight decreased	6 (2.0)
Aspartate aminotransferase increased	4 (1.3)
<b>METABOLISM AND NUTRITION DISORDERS</b>	
Hypophosphatemia	17 (5.6)



**Table 3. Percentage of Patients with Adverse Events  $\geq$  1% Reported in Patients with Giant Cell Tumour of Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N=304) n (%)</b>
Hypocalcemia	13 (4.3)
Decreased appetite	11 (3.6)
Hypercalcemia	6 (2.0)
Hyperglycemia	5 (1.6)
Hypokalemia	4 (1.3)
<b>MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS</b>	
Arthralgia	64 (21.1)
Back pain	53 (17.4)
Pain in extremity	49 (16.1)
Musculoskeletal pain	26 (8.6)
Muscle spasms	17 (5.6)
Bone pain	16 (5.3)
Myalgia	16 (5.3)
Neck pain	15 (4.9)
Pain in jaw	10 (3.3)
Muscular weakness	6 (2.0)
Musculoskeletal chest pain	5 (1.6)
Musculoskeletal stiffness	5 (1.6)
Joint swelling	4 (1.3)
Osteonecrosis of jaw	4 (1.3)
<b>NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS)</b>	
Tumour pain	9 (3.0)
<b>NERVOUS SYSTEM DISORDERS</b>	
Headache	56 (18.4)
Paresthesia	16 (5.3)
Dizziness	15 (4.9)
Hypoesthesia	10 (3.3)
Neuralgia	8 (2.6)
Lethargy	4 (1.3)
Peripheral sensory neuropathy	4 (1.3)
Somnolence	4 (1.3)
<b>PSYCHIATRIC DISORDERS</b>	
Insomnia	16 (5.3)
Depression	14 (4.6)
Anxiety	8 (2.6)
<b>RENAL AND URINARY DISORDERS</b>	
Dysuria	4 (1.3)
Urinary incontinence	4 (1.3)

**Table 3. Percentage of Patients with Adverse Events  $\geq$  1% Reported in Patients with Giant Cell Tumour of Bone by System Organ Class**

<b>SYSTEM ORGAN CLASS Preferred Term</b>	<b>XGEVA (N=304) n (%)</b>
<b>REPRODUCTIVE SYSTEM AND BREAST DISORDERS</b>	
Pelvic pain	6 (2.0)
<b>RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS</b>	
Cough	19 (6.3)
Dyspnea	11 (3.6)
Oropharyngeal pain	10 (3.3)
Nasal congestion	6 (2.0)
Rhinorrhea	4 (1.3)
<b>SKIN AND SUBCUTANEOUS TISSUE DISORDERS</b>	
Rash	15 (4.9)
Pruritus	9 (3.0)
Eczema	7 (2.3)
Acne	6 (2.0)
Dry skin	6 (2.0)
Alopecia	5 (1.6)
<b>VASCULAR DISORDERS</b>	
Hot flush	12 (3.9)
Hypertension	4 (1.3)

N = Number of subjects who received  $\geq$  1 active dose of investigational product

n = Number of subjects reporting  $\geq$  1 event

Includes only treatment-emergent adverse events

System organ classes are sorted alphabetically and preferred terms are sorted by descending order of frequency and coded using MedDRA version 14.1

Subjects who rolled over from Study 5 to Study 6 or who discontinued Study 5 and re-entered Study 6 are counted only once and their analysis period will start from Study 5 and end at Study 6

In the final analysis of Study 6, of the 526 subjects who received  $\geq$  1 dose of XGEVA, the most common treatment-emergent adverse events were arthralgia (35.0%), back pain (28.3%), fatigue (27.0%), pain in extremity (27.0%), headache (25.1%), nausea (22.1%), nasopharyngitis (14.4%), toothache (12.9%), musculoskeletal pain (12.7%), hypophosphatemia (12.4%), vomiting (12.2%), diarrhea (11.2%), constipation (11.0%) and cough (10.3%). The most common treatment-emergent serious adverse events were osteonecrosis of the jaw (3.8%), bone giant cell tumour (1.5%), anemia (1.1%), and back pain (1.0%). In addition, the following treatment-emergent adverse events were also observed, musculoskeletal pain (68.4%), infections (53.4%), adverse events potentially associated with hypersensitivity (20.2%), vascular disorders (10.3%) and cardiac disorders (6.1%). The incidence of treatment-emergent fatal adverse events was 2.1% (11 of 526 subjects) with three deaths being considered by investigators as being related to denosumab (0.6%; bone sarcoma, rhabdomyosarcoma and

sarcoma). The incidence of treatment-emergent adverse events leading to treatment phase discontinuation was 9.9% (52 of 526 patients).

### Hypercalcemia of Malignancy Refractory to Intravenous Bisphosphonate

The safety of XGEVA was evaluated in an open-label, single-arm trial (Study 7) in which 33 patients were enrolled with hypercalcemia of malignancy (with or without bone metastases) refractory to treatment with intravenous bisphosphonate. Patients received XGEVA subcutaneously every 4 weeks with additional 120 mg doses on Days 8 and 15 of the first month of therapy.

Of the 33 patients who received XGEVA, 33 patients were treated with XGEVA for  $\geq 1$  month, 5 patients for  $\geq 6$  months, and 3 patients for  $\geq 1$  year. The median number of doses received was 4 (range: 1 to 25 doses) and the median number of months on study was 1.8 (range: 0 to 23 months). Sixty-four percent of enrolled patients were men and 70% were White. The median age was 63 years (range: 22 to 89 years).

The adverse event profile of XGEVA in patients with hypercalcemia of malignancy was similar to that reported in Studies 1, 2, 3, 5, and 6 (Table 4). The most common adverse events were nausea (10 patients, 30%), dyspnea (9 patients, 27%), and decreased appetite, headache, peripheral edema, and vomiting, each of which occurred in 8 patients (24%). The patient incidence of fatal adverse events was 78.8%. The most frequently reported serious adverse events were worsening hypercalcemia (5 subjects, 15%), dyspnea (3 subjects, 9%), and metastatic breast cancer (3 subjects, 9%). No adverse events leading to discontinuation were reported as related to XGEVA treatment.

**Table 4. Treatment-Emergent Adverse Events by Preferred Term in Descending Order of Frequency ( $\geq 5\%$  Patient Incidence)**

Preferred Term	XGEVA (N = 33) n (%)
Number of patients reporting adverse events	32 (97.0)
Nausea	10 (30.3)
Dyspnea	9 (27.3)
Decreased appetite	8 (24.2)
Headache	8 (24.2)
Edema peripheral	8 (24.2)
Vomiting	8 (24.2)
Anemia	7 (21.2)
Constipation	7 (21.2)
Diarrhea	7 (21.2)
Cough	6 (18.2)
Fatigue	6 (18.2)
Hypercalcemia	6 (18.2)
Hypotension	5 (15.2)
Pleural effusion	5 (15.2)
Pyrexia	5 (15.2)
Arthralgia	4 (12.1)
Back pain	4 (12.1)
Confusional state	4 (12.1)

**Table 4. Treatment-Emergent Adverse Events by Preferred Term in Descending Order of Frequency (≥ 5% Patient Incidence)**

<b>Preferred Term</b>	<b>XGEVA (N = 33) n (%)</b>
Fluid overload	4 (12.1)
Hypophosphatemia	4 (12.1)
Pulmonary Edema	4 (12.1)
Weight decreased	4 (12.1)
Abdominal distension	3 (9.1)
Abdominal pain	3 (9.1)
Alopecia	3 (9.1)
Breast cancer metastatic	3 (9.1)
Epistaxis	3 (9.1)
Hemoglobin decreased	3 (9.1)
Hypocalcemia	3 (9.1)
Hypokalemia	3 (9.1)
Hyponatremia	3 (9.1)
Myalgia	3 (9.1)
Oropharyngeal pain	3 (9.1)
Pneumonia	3 (9.1)
Rash	3 (9.1)
Renal failure acute	3 (9.1)
Tachycardia	3 (9.1)
Thrombocytopenia	3 (9.1)
Asthenia	2 (6.1)
Breast cancer	2 (6.1)
<i>Clostridium difficile</i> colitis	2 (6.1)
Dehydration	2 (6.1)
Depressed level of consciousness	2 (6.1)
Dizziness	2 (6.1)
Dry mouth	2 (6.1)
Dysphonia	2 (6.1)
Fall	2 (6.1)
Febrile neutropenia	2 (6.1)
Head and neck cancer	2 (6.1)
Hyperkalemia	2 (6.1)
Hypomagnesemia	2 (6.1)
Hypoxia	2 (6.1)
Insomnia	2 (6.1)
Lethargy	2 (6.1)
Lung neoplasm malignant	2 (6.1)
Musculoskeletal pain	2 (6.1)
Neutropenia	2 (6.1)
Non-small cell lung cancer	2 (6.1)
Pain in extremity	2 (6.1)
Palmar-plantar erythrodysesthesia syndrome	2 (6.1)
Plasma cell myeloma	2 (6.1)

**Table 4. Treatment-Emergent Adverse Events by Preferred Term in Descending Order of Frequency (≥ 5% Patient Incidence)**

<b>Preferred Term</b>	<b>XGEVA (N = 33) n (%)</b>
Rash pruritic	2 (6.1)
Rhinorrhea	2 (6.1)

N = Number of patients who received ≥ 1 dose of XGEVA; n = Number of patients reporting ≥ 1 adverse event; Includes only treatment-emergent adverse events; Preferred terms are sorted by descending order of frequency and coded using MedDRA version 16.0

**Table 5. Investigational Product Related Adverse Events by Preferred Term in Descending Order of Frequency**

<b>Preferred Term</b>	<b>XGEVA (N = 33) n (%)</b>
Number of subjects reporting investigational product related adverse events	13 (39.4)
Hypophosphatemia	4 (12.1)
Nausea	4 (12.1)
Edema peripheral	3 (9.1)
Arthralgia	2 (6.1)
Constipation	2 (6.1)
Diarrhea	2 (6.1)
Hypocalcemia	2 (6.1)
Myalgia	2 (6.1)
Pneumonia	2 (6.1)
Rash	2 (6.1)
Rash pruritic	2 (6.1)
Rhinorrhea	2 (6.1)
Vomiting	2 (6.1)
Abdominal distension	1 (3.0)
Abdominal pain	1 (3.0)
Abdominal pain lower	1 (3.0)
Aspartate aminotransferase increased	1 (3.0)
Back pain	1 (3.0)
Bone pain	1 (3.0)
Cardiac arrest	1 (3.0)
Cardiac murmur	1 (3.0)
<i>Clostridium difficile</i> colitis	1 (3.0)
Coccydynia	1 (3.0)
Colitis	1 (3.0)
Contusion	1 (3.0)
Dysgeusia	1 (3.0)
Dyspepsia	1 (3.0)
Electrocardiogram QT prolonged	1 (3.0)
Fatigue	1 (3.0)

**Table 5. Investigational Product Related Adverse Events by Preferred Term in Descending Order of Frequency**

<b>Preferred Term</b>	<b>XGEVA (N = 33) n (%)</b>
Fluid overload	1 (3.0)
Headache	1 (3.0)
Hypertension	1 (3.0)
Lung infection	1 (3.0)
Musculoskeletal pain	1 (3.0)
Musculoskeletal stiffness	1 (3.0)
Oropharyngeal pain	1 (3.0)
Pharyngitis	1 (3.0)
Pollakiuria	1 (3.0)
Respiratory tract infection	1 (3.0)
Temporomandibular joint syndrome	1 (3.0)
Toothache	1 (3.0)
Urinary retention	1 (3.0)
Ventricular tachycardia	1 (3.0)
Vision blurred	1 (3.0)

N = Number of subjects who received  $\geq 1$  dose of investigational product

n = Number of subjects reporting  $\geq 1$  event

Includes only treatment-emergent adverse events for which the investigator indicated there was a reasonable possibility they may have been caused by investigational product

Preferred terms are sorted by descending order of frequency and coded using MedDRA version 15.1

One fatal event (cardiac arrest) and 1 serious grade 4 colitis were reported by the investigator to be related to XGEVA therapy. The XGEVA therapy-related grade 3 events of fatigue, infections and hypophosphatemia were reported in 3%, 6.1% and 9.1% of the patients, respectively.

### **Hypocalcemia**

In clinical trials in patients with advanced cancer and adequate renal function (defined as estimated creatinine clearance  $\geq 30$  mL/min), hypocalcemia was reported as an adverse event in 9.6% of patients in the XGEVA group and 5.0% of patients in the zoledronic acid group.

Severe hypocalcemia (corrected serum calcium less than 7 mg/dL or less than 1.75 mmol/L) occurred in 3.1% of patients treated with XGEVA and 1.3% of patients treated with zoledronic acid. Of patients who experienced severe hypocalcemia, 33% experienced 2 or more episodes of severe hypocalcemia and 16% experienced 3 or more episodes (see **7 WARNINGS AND PRECAUTIONS, Endocrine and Metabolism, Hypocalcemia**).

Initially in clinical trials in patients with GCTB, moderate hypocalcemia (corrected serum calcium less than 8 to 7 mg/dL or less than 2 to 1.75 mmol/L) occurred in 2.6% of patients treated with XGEVA. At the final integrated analysis of clinical trials in patients with GCTB, moderate hypocalcemia (as defined above) occurred in 4.4% of patients treated with XGEVA.

Two clinical trials were conducted in subjects without cancer and with varying degrees of renal function.

In one study, subjects (N = 55) with varying degrees of renal function (ranging from normal through end-stage renal disease requiring dialysis) received a single 60 mg subcutaneous dose of denosumab. Hypocalcemia was observed in 8 subjects (15%), 1 (2%) of whom was symptomatic. Two subjects (4%) each experienced an adverse event of hypocalcemia that was

classified as serious. Both subjects had severe chronic kidney disease (CKD) and were enrolled before the protocol required supplementation of calcium and vitamin D.

In a second study, patients (N = 32) with severe renal impairment (creatinine clearance less than 30 mL/minute and/or on dialysis) were given two 120 mg subcutaneous doses (Days 1 and 29) of denosumab. Two patients overall (1 in each group) had symptomatic hypocalcemia, based on clinical adverse events and concomitant symptoms. One patient in the severe CKD group had concomitant muscle spasms and 1 patient in the CKD on dialysis group had concomitant paresthesia. Hence, the overall incidence of clinically significant hypocalcemia (corrected serum calcium less than 1.75 mmol/L or symptomatic hypocalcemia) was 9.4%: 1 of 16 patients (6.3%) in the severe group and 2 of 16 patients (12.5%) in the CKD on dialysis group. Both events of symptomatic hypocalcemia were mild in severity. Accompanying increases in parathyroid hormone (PTH) have also been observed in patients receiving XGEVA with severe renal impairment or receiving dialysis. At baseline, median (range) intact PTH (iPTH) values were 6.1 pmol/L (1.06 pmol/L, 13.2 pmol/L) and 16.3 pmol/L (0.3 pmol/L, 37.0 pg/mL) in the severe CKD and CKD on dialysis groups, respectively. At end of study (Day 113), median (range) iPTH values were 7.1 pmol/L (0.85 pmol/L, 372.2 pmol/L) and 31.5 pmol/L (1.3 pmol/L, 136.1 pmol/mL) in the severe CKD and CKD on dialysis groups, respectively. The median percent change from baseline to end of study (Day 113) was 15.0% in the severe CKD group and 107.5% in the CKD on dialysis group. The median (range) maximal elevation of PTH was 19.4 pmol/L (2.8 pmol/L, 372.2 pmol/L) in the severe CKD group and 64.2 pmol/L (1.3 pmol/L, 396.1 pmol/L) in the CKD on dialysis group. The median (range) % change of the maximal elevation of PTH values from baseline was 164.8% (28.2%, 2729.0%) in the severe CKD group and 256.1% (46.8%, 1681.9%) in the CKD on dialysis group. The median (range) time to reach maximal elevation of PTH was 22.5 days (3 days, 115 days) in the severe CKD group and 64.0 days (6 days, 115 days) in the CKD dialysis group. The median duration (range) of the elevation (20% above baseline) of PTH was 41.0 days (3.0 days, 112.0 days) in the severe CKD group and 79.5 days (11.0 days, 216.0 days) in the CKD on dialysis group.

In both studies, there was a greater risk of developing hypocalcemia with increasing degree of renal impairment, and in the absence of, or inadequate calcium supplementation.

### **Osteonecrosis of the Jaw (ONJ)**

In clinical trials in patients with advanced cancer, ONJ was confirmed in 1.8% of patients in the XGEVA group (median exposure of 12.0 months; range 0.1 – 40.5) and 1.3% of patients in the zoledronic acid group (see **7 WARNINGS AND PRECAUTIONS**). Fifty-eight percent of subjects in the XGEVA group and 65% of subjects in the zoledronic acid group had a prior or concurrent tooth extraction, 42% of subjects in the XGEVA group and 27% of subjects in the zoledronic acid group had used a denture or other dental appliance, and 31% of subjects in the XGEVA group and 32% of subjects in the zoledronic acid group had poor oral hygiene.

The trials in patients with breast or prostate cancer included an XGEVA extension treatment phase (median overall exposure of 14.9 months; range 0.1 – 67.2) (see **14 CLINICAL TRIALS**). For patients who were randomized to XGEVA and continued on XGEVA in the open label extension phase, the patient-year adjusted incidence of confirmed ONJ was 1.1 per 100 patient-years during the first year of treatment, 3.7 in the second year, and 4.6 thereafter. The median time to ONJ was 20.6 months (range: 4 – 53) (see **7 WARNINGS AND PRECAUTIONS**).

In a phase 3 double-blind, active-controlled clinical trial in patients with newly diagnosed multiple myeloma, ONJ was confirmed in 4.1% of patients in the XGEVA group (median exposure of 15.8 months; range 1 – 49.8) and 2.8% of patients in the zoledronic acid group. At the completion of the double-blind treatment phase of this trial, the patient-year adjusted

incidence of confirmed ONJ in the XGEVA group (median exposure of 19.4 months; range 1 – 52) was 2.0 per 100 patient-years during the first year of treatment, 5.0 in the second year, and 4.5 thereafter. The median time to ONJ was 18.7 months (range: 1 - 44) (see **7 WARNINGS AND PRECAUTIONS**).

In clinical trials in patients with GCTB, at initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6), ONJ was confirmed in 4 of 304 (1.3%) patients who received XGEVA. The median time to ONJ was 16 months (range: 13 to 20 months).

A final analysis of a phase 2 open-label clinical trial (Study 6) in patients with GCTB demonstrated that ONJ was confirmed in 36 of 526 (6.8%) patients (median number of 34 doses; range 4 – 116). Of these 36 cases, 26 (4.9%) were reported as serious adverse events by investigators. At the completion of the trial, median time on trial including safety follow-up phase was 60.9 months (range: 0 – 112.6). The patient-year adjusted incidence of confirmed ONJ was 1.5 per 100 patient-years overall (0.2 per 100 patient-years during the first year of treatment, 1.5 in the second year, 1.8 in the third year, 2.1 in the fourth year, 1.4 in the fifth year, and 2.2 thereafter). The median time to ONJ was 41 months (range: 11 – 96) (see **7 WARNINGS AND PRECAUTIONS**).

### **Atypical Femoral Fractures**

In the clinical trial program, atypical femoral fracture has been reported uncommonly in patients treated with XGEVA 120 mg and the risk increased with longer duration of treatment. Events have occurred during treatment and up to 9 months after treatment was discontinued.

### **Malignancies**

In a pooled safety analysis of clinical trials in cancer patients with bone metastases, the overall incidence of new primary malignancies was 0.99% (28 out of 2841 patients) in the XGEVA group and 0.63% (18 out of 2836 patients) in the zoledronic acid group. In the breast cancer trial, the incidence was 0.5 % in both XGEVA (5/1020 patients) and zoledronic acid groups (5/1013 patients). In other solid tumours or multiple myeloma, the incidence was 0.6% (5/878 patients) and 0.3% (3/878 patients) in the XGEVA and zoledronic acid groups, respectively. In the prostate cancer trial, the incidence was 1.9% (18/943 patients) in the XGEVA group and 1.1% (10/945 patients) in the zoledronic acid group. In the final analysis of the GCTB trial (Study 6), the incidence of malignancy was 3.8% (20/526 patients, all adults).



### 8.3 Less Common Clinical Trial Adverse Reactions

#### Less Common Clinical Trial Adverse Events (< 1%) in Patients with Advanced Malignancies Involving Bone by System Organ Class

**BLOOD AND LYMPHATIC SYSTEM DISORDERS:** leukocytosis, lymphadenopathy, lymphopenia, coagulopathy, neutrophilia, disseminated intravascular coagulation, thrombocytosis, splenomegaly, lymph node pain, hemorrhagic diathesis, anemia of chronic disease, lymphadenopathy mediastinal, macrocytosis, iron deficiency anemia

**CARDIAC DISORDERS:** arrhythmia, pericardial effusion, cardiac failure congestive, angina pectoris, cardiac arrest, cardio-respiratory arrest, myocardial ischemia, cardiopulmonary failure, myocardial infarction, sinus tachycardia, acute myocardial infarction, cardiac failure acute, mitral valve incompetence, pericarditis, coronary artery disease, left ventricular hypertrophy, supraventricular tachycardia, cardiomyopathy, cardiogenic shock, bradycardia, left ventricular failure, tricuspid valve incompetence, angina unstable, bundle branch block right, ventricular tachycardia, cardiomegaly, acute coronary syndrome, extrasystoles, atrial flutter, cardiovascular insufficiency, cardiovascular disorder, diastolic dysfunction

**CONGENITAL, FAMILIAL AND GENETIC DISORDERS:** phimosis

**EAR AND LABYRINTH DISORDERS:** ear pain, tinnitus, hypoacusis, hearing impaired, deafness, cerumen impaction, ear pruritus, vertigo positional, ear discomfort

**ENDOCRINE DISORDERS:** cushingoid, hypothyroidism, goiter, hyperthyroidism, cushing's syndrome, adrenal insufficiency

**EYE DISORDERS:** visual acuity reduced, visual impairment, diplopia, dry eye, cataract, eye pain, eye irritation, eye hemorrhage, eyelid ptosis, eye pruritus, conjunctival hemorrhage, eyelid edema, eye swelling, myodesopsia, exophthalmos, blepharitis, eye inflammation, photophobia, photopsia, conjunctivitis allergic, keratoconjunctivitis sicca, retinal detachment, blindness unilateral, eye disorder, keratitis, ophthalmoplegia, eye edema, ocular hyperemia, glaucoma, amaurosis, lacrimal disorder

**GASTROINTESTINAL DISORDERS:** abdominal pain lower, esophagitis, dental caries, oral pain, mouth ulceration, periodontitis, ileus, tooth disorder, hematochezia, loose tooth, hypoesthesia oral, odynophagia, intestinal obstruction, aphthous stomatitis, gastrointestinal hemorrhage, hematemesis, proctalgia, colitis, gingival bleeding, hiatus hernia, periodontal disease, diverticulum, hemorrhoidal hemorrhage, gingival recession, sensitivity of teeth, paresthesia oral, melena, glossodynia, epigastric discomfort, tooth loss, upper gastrointestinal hemorrhage, gastrointestinal disorder, cheilitis, irritable bowel syndrome, small intestinal obstruction, mouth hemorrhage, fecal incontinence, duodenal ulcer, inguinal hernia, hyperchlorhydria, gastric ulcer, oral discomfort, salivary hypersecretion, lip dry, lip swelling, lower gastrointestinal hemorrhage, oral disorder, retching, gingival ulceration, gastritis erosive, duodenitis, gastrointestinal obstruction, anal hemorrhage, fecaloma, abdominal tenderness, peritonitis, gastrointestinal motility disorder, gingival disorder, subileus, gastrointestinal edema, anal fissure, feces discoloured, diverticulum intestinal, eructation, gingival swelling, intestinal perforation, esophageal stenosis, colonic polyp, proctitis, umbilical hernia, anal pruritus, gingival erythema, edema mouth, rectal tenesmus, reflux gastritis, frequent bowel movements, gastric ulcer hemorrhage, obstruction gastric, oral dysesthesia, reflux esophagitis, anorectal discomfort, gastrointestinal hypomotility, gastrointestinal pain, enteritis, tongue discoloration, tongue disorder

**GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS:** performance status decreased, death, local swelling, catheter related complication, hyperthermia, axillary pain,

generalized edema, localized edema, facial pain, catheter site pain, injection site pain, injection site reaction, feeling cold, inflammation, adverse drug reaction, extravasation, non-cardiac chest pain, xerosis, temperature intolerance, thirst, early satiety, sudden death, nodule, suprapubic pain, abasia, impaired healing, injection site pruritus, cyst, irritability, catheter site hematoma, infusion site pain, catheter site inflammation, feeling hot, swelling, catheter site hemorrhage, hernia, catheter site erythema, catheter thrombosis, injection site hematoma, drug withdrawal syndrome, infusion site extravasation, discomfort, injection site hemorrhage, drug intolerance, hypothermia, mucosal dryness, organ failure, induration

**HEPATOBIILIARY DISORDERS:** hyperbilirubinemia, hepatomegaly, cholelithiasis, hepatic pain, hepatic steatosis, liver disorder, cholecystitis, cholestasis, hepatitis toxic, hepatorenal failure, bile duct obstruction, hepatic cyst, hepatotoxicity, hepatic lesion

**IMMUNE SYSTEM DISORDERS:** hypersensitivity, drug hypersensitivity, seasonal allergy

**INFECTIONS AND INFESTATIONS:** tooth infection, localized infection, lung infection, catheter related infection, infection, erysipelas, candidiasis, paronychia, oral fungal infection, septic shock, respiratory tract infection viral, viral infection, lobar pneumonia, skin infection, onychomycosis, nail infection, urosepsis, diverticulitis, wound infection, ear infection, herpes virus infection, eye infection, vaginal infection, pyelonephritis, subcutaneous abscess, laryngitis, bronchopneumonia, osteomyelitis, gastroenteritis viral, fungal skin infection, oral infection, tinea pedis, catheter site infection, gingival infection, vulvovaginal mycotic infection, furuncle, neutropenic sepsis, esophageal candidiasis, fungal infection, tracheitis, clostridial infection, tonsillitis, postoperative wound infection, *Clostridium difficile* colitis, Staphylococcal infection, viral upper respiratory tract infection, breast cellulitis, staphylococcal sepsis, gastrointestinal infection, bacteremia, infected bites, lymphangitis, folliculitis, gingival abscess, otitis media, vulvovaginal candidiasis, abscess limb, *Escherichia* bacteremia, lung abscess, pyelonephritis acute, hordeolum, infected skin ulcer, labyrinthitis, mastitis, orchitis, soft tissue infection, gangrene, genital infection fungal, pyelonephritis chronic, acarodermatitis, pulpitis dental, febrile infection, kidney infection, skin candida, herpes simplex, abscess, catheter site cellulitis, Enterococcal infection, pneumonia *Klebsiella*

**INJURY, POISONING AND PROCEDURAL COMPLICATIONS:** procedural pain, radiation skin injury, cervical vertebral fracture, tooth fracture, wound, pelvic fracture, fractured ischium, fracture, ilium fracture, skin laceration, fractured sacrum, limb injury, humerus fracture, muscle strain, thermal burn, clavicle fracture, subdural hematoma, excoriation, joint sprain, radius fracture, drug toxicity, head injury, joint injury, arthropod bite, scapula fracture, tooth injury, tibia fracture, fibula fracture, gastroenteritis radiation, medical device complication, post-traumatic pain, sternal fracture, wound complication, post procedural hemorrhage, radiation pneumonitis, sunburn, contrast media reaction, ulna fracture, seroma, device breakage, transfusion reaction, eye injury, radiation injury, skeletal injury, subdural hemorrhage, concussion, post procedural complication, wound dehiscence, face injury, joint dislocation, poisoning, tendon rupture, animal bite, facial bones fracture, overdose, radiation associated pain

**INVESTIGATIONS:** blood bilirubin increased, white blood cell count decreased, gamma-glutamyltransferase increased, blood alkaline phosphatase, body temperature increased, blood calcium decreased, blood urea increased, blood pressure increased, blood glucose increased, blood potassium increased, blood potassium decreased, international normalized ratio increased, neutrophil count decreased, hemoglobin, blood albumin decreased, hepatic enzyme increased, liver function test abnormal, blood lactate dehydrogenase increased, blood phosphorus decreased, hematocrit decreased, white blood cell count increased, blood magnesium decreased, blood bicarbonate decreased, blood creatinine, urine output decreased, cardiac murmur, red blood cell count decreased, blood sodium decreased, blood uric acid

increased, blood urine present, blood iron decreased, transaminases increased, blood creatine increased, heart rate increased, neutrophil count, platelet count increased, creatinine renal clearance decreased, blood creatinine decreased, bone density decreased, electrocardiogram QT prolonged, occult blood positive, protein total decreased, C-reactive protein increased, breath sounds abnormal, prothrombin time prolonged, activated partial thromboplastin time prolonged, blood calcium increased, Eastern Cooperative Oncology Group performance status worsened, ejection fraction decreased, neutrophil count increased

**METABOLISM AND NUTRITION DISORDERS:** diabetes mellitus, malnutrition, hypercholesterolemia, hyperuricemia, hypoproteinemia, metabolic acidosis, gout, fluid retention, hypercreatininemia, failure to thrive, hypertriglyceridemia, electrolyte imbalance, dyslipidemia, hypermagnesemia, hypophagia, hypovolemia, fluid overload, iron deficiency, hyperlipidemia, diabetes mellitus inadequate control, type 2 diabetes mellitus, acidosis, vitamin D deficiency, polydipsia, tumour lysis syndrome, appetite disorder, vitamin B12 deficiency

**MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS:** osteoarthritis, musculoskeletal discomfort, joint stiffness, joint range of motion decreased, arthritis, coccydynia, pubic pain, intervertebral disc protrusion, bone lesion, limb discomfort, spinal osteoarthritis, tendonitis, myopathy, mobility decreased, muscle tightness, jaw disorder, rotator cuff syndrome, osteolysis, bursitis, peri-arthritis, sensation of heaviness, hypercreatininemia, muscle fatigue, osteoporosis, osteitis, tendon disorder, exostosis, amyotrophy, intervertebral disc degeneration, dupuytren's contracture, rheumatoid arthritis, trigger finger, arthropathy, muscle atrophy, joint effusion, osteopenia, pathological fracture, plantar fasciitis, muscle twitching, spinal disorder, spondylolisthesis, scoliosis, muscle contracture, joint crepitation, lumbar spinal stenosis, muscle rigidity, tenosynovitis, trismus

**NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS):** metastasis, cancer pain, breast cancer, metastases to lymph nodes, benign neoplasm of skin, prostate cancer metastatic, metastatic pain, malignant pleural effusion, metastases to meninges, breast cancer metastatic, metastases to bone marrow, metastatic neoplasm, lung cancer metastatic, basal cell carcinoma, malignant ascites, seborrheic keratosis, tumour pain, metastases to pleura, benign neoplasm, paraneoplastic syndrome, skin papilloma, metastases to peritoneum, colon cancer metastatic, gastric cancer, metastases to bladder, rectal cancer, benign neoplasm of thyroid gland, meningioma, metastases to gastrointestinal tract, metastases to skin, neoplasm progression, squamous cell carcinoma, lip neoplasm benign, tumour associated fever

**NERVOUS SYSTEM DISORDERS:** cerebrovascular accident, peripheral motor neuropathy, balance disorder, facial palsy, memory impairment, ataxia, polyneuropathy, amnesia, dysarthria, loss of consciousness, dysesthesia, monoparesis, migraine, paraparesis, hyperesthesia, neurotoxicity, hemiparesis, presyncope, cerebral ischemia, brain edema, intracranial pressure increased, transient ischemic attack, cervical cord compression, speech disorder, hemiplegia, cognitive disorder, sensory disturbance, disturbance in attention, ageusia, vocal cord paralysis, carpal tunnel syndrome, coma, epilepsy, restless legs syndrome, sinus headache, hypertonia, burning sensation, aphasia, paraplegia, hypotonia, aphonia, monoplegia, dizziness postural, motor dysfunction, cerebral hemorrhage, hydrocephalus, dementia, encephalopathy, paresis, parosmia, paralysis, poor quality sleep, depressed level of consciousness, hepatic encephalopathy, ischemic stroke, cerebrovascular disorder, cervical root pain, cranial neuropathy, facial paresis, epiduritis, grand mal convulsion, sensory loss, autonomic nervous system imbalance, cauda equina syndrome, cerebellar syndrome, neurological decompensation, partial seizures, cerebral atrophy, cerebral infarction, trigeminal neuralgia, hypogeusia, mental impairment, nervous system disorder, vascular encephalopathy, peroneal

*nerve palsy, hemorrhage intracranial, horner's syndrome, coordination abnormal, dyskinesia, hypersomnia, formication, hemorrhagic stroke, radiculopathy, sedation, radicular pain, diplegia*

**PSYCHIATRIC DISORDERS:** *sleep disorder, disorientation, mental status changes, depressed mood, hallucination, mood altered, restlessness, nervousness, delirium, affect lability, abnormal behaviour, libido decreased, nightmare, bruxism, neurosis, adjustment disorder, fear, mental disorder, panic attack, mood swings, apathy, stress, hallucination (visual), aggression*

**RENAL AND URINARY DISORDERS:** *urinary tract obstruction, urinary tract disorder, oliguria, renal cyst, nephrolithiasis, proteinuria, micturition urgency, incontinence, renal pain, renal failure chronic, urinary bladder hemorrhage, micturition disorder, urinary hesitation, urine flow decreased, bladder spasm, bladder obstruction, polyuria, anuria, hemorrhage urinary tract, ureteric obstruction, calculus bladder, bladder pain, renal colic, calculus ureteric, urethral stenosis, hypertonic bladder, urethral obstruction, bladder disorder, choluria, azotemia, chromaturia, obstructive uropathy, renal disorder, bladder neck obstruction, strangury*

**REPRODUCTIVE SYSTEM AND BREAST DISORDERS:** *gynecomastia, vaginal hemorrhage, vulvovaginal dryness, scrotal edema, nipple pain, penile edema, edema genital, vaginal discharge, benign prostatic hyperplasia, erectile dysfunction, ovarian cyst, prostatitis, vulvovaginal pruritus, amenorrhea, prostatism, scrotal pain, metrorrhagia, genital hemorrhage, breast edema, breast mass, breast tenderness, balanitis, breast discomfort, dyspareunia, menstruation irregular, vaginal inflammation, vulvovaginal pain, perineal pain, genital discharge, penile pain, testicular pain, breast hemorrhage, breast swelling, pruritus genital, menorrhagia, pelvic discomfort*

**RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS:** *wheezing, hydrothorax, atelectasis, chronic obstructive pulmonary disease, hiccups, rhinitis allergic, lung infiltration, pleurisy, pulmonary edema, pleuritic pain, bronchospasm, sinus congestion, respiratory tract congestion, lung disorder, pneumothorax, acute respiratory failure, postnasal drip, respiratory distress, rhonchi, pneumonitis, rales, acute respiratory distress syndrome, bronchitis chronic, nasal dryness, asthma, respiratory disorder, tachypnea, dry throat, increased bronchial secretion, pulmonary hypertension, paranasal sinus hypersecretion, orthopnea, emphysema, sinus disorder, hydropneumothorax, hemothorax, throat irritation, increased upper airway secretion, pulmonary hemorrhage, bronchial obstruction, asphyxia, nasal discomfort, pharyngeal inflammation, respiratory tract hemorrhage, apnea, pulmonary fibrosis, respiratory arrest, upper airway obstruction, acute pulmonary edema, hypoventilation, pulmonary congestion, interstitial lung disease, aspiration*

**SKIN AND SUBCUTANEOUS TISSUE DISORDERS:** *skin lesion, hypoesthesia facial, decubitus ulcer, urticaria, ecchymosis, swelling face, skin reaction, skin hyperpigmentation, skin exfoliation, dermatitis allergic, acne, onycholysis, skin discolouration, rash erythematous, eczema, nail discolouration, blister, petechiae, skin disorder, periorbital edema, dermatitis contact, skin nodule, subcutaneous nodule, skin irritation, onychoclasis, rash popular, rash pruritic, skin fissures, ingrowing nail, rash macular, actinic keratosis, pain of skin, dermatitis acneiform, pigmentation disorder, pruritus generalized, nail toxicity, drug eruption, seborrheic dermatitis, cold sweat, hyperkeratosis, rash generalized, skin atrophy, skin edema, xeroderma, skin toxicity, exfoliative rash, nail dystrophy, dermal cyst, skin hemorrhage, purpura, intertrigo, onychomadesis, increased tendency to bruise, psoriasis, photodermatitis, ingrown hair, scar*

**SOCIAL CIRCUMSTANCES:** *immobile*

**SURGICAL AND MEDICAL PROCEDURES:** *mastectomy, catheter placement, endodontic procedure, bladder catheterization, cataract operation, transurethral prostatectomy, central venous catheterization*

**VASCULAR DISORDERS:** hematoma, thrombosis, flushing, thrombophlebitis, pallor, hemorrhage, hypertensive crisis, lymphostasis, orthostatic hypotension, peripheral coldness, venous thrombosis, arteriosclerosis, varicose vein, circulatory collapse, venous thrombosis limb, jugular vein thrombosis, superior vena caval occlusion, aortic aneurysm, aortic arteriosclerosis, venous insufficiency, subclavian vein thrombosis, intermittent claudication, phlebitis superficial, hypovolemic shock, thrombophlebitis superficial, vasculitis, embolism, vein pain

**Less Common Clinical Trial Adverse Events\* (< 1%) in Patients with Giant Cell Tumour of Bone by System Organ Class**

**BLOOD AND LYMPHATIC SYSTEM DISORDERS:** leukopenia, lymphadenopathy, hypochromic anemia, iron deficiency anemia, lymph node pain, spleen disorder, splenomegaly

**CARDIAC DISORDERS:** sinus tachycardia, pericarditis

**EAR AND LABYRINTH DISORDERS:** ear pain, eustachian tube obstruction, tympanic membrane perforation

**ENDOCRINE DISORDERS:** hyperthyroidism, goitre, hyperparathyroidism, hypothyroidism, toxic nodular goitre

**EYE DISORDERS:** cataract, dry eye, eye irritation, visual impairment, abnormal sensation in eye, astigmatism, diplopia, eye edema, eye pruritus, lacrimation increased, myopia, ocular hyperemia, periorbital edema, photophobia, visual acuity reduced

**GASTROINTESTINAL DISORDERS:** abdominal pain lower, flatulence, hematochezia, rectal hemorrhage, sensitivity of teeth, abdominal rigidity, aphthous stomatitis, chapped lips, cheilitis, gingival bleeding, gingival disorder, gingivitis, lip ulceration, odynophagia, oral pain, pancreas lipomatosis, periodontitis, tooth discoloration, umbilical hernia

**GENERAL DISORDERS AND ADMINISTRATION SITE CONDITIONS:** feeling cold, edema, atrophy, axillary pain, cyst, device failure, gait disturbance, hyperthermia, injection site erythema, injection site irritation, localized edema, mucosal inflammation, nodule, sensation of pressure, temperature intolerance

**HEPATOBIILIARY DISORDERS:** cholelithiasis, hepatic steatosis

**IMMUNE SYSTEM DISORDERS:** allergy to animal, hypersensitivity, seasonal allergy

**INFECTIONS AND INFESTATIONS:** oral herpes, osteomyelitis, skin infection, viral upper respiratory tract infection, abscess oral, acarodermatitis, acute tonsillitis, appendicitis, dental fistula, device related infection, eye infection, folliculitis, gastritis bacterial, genital candidiasis, genital herpes, helicobacter infection, infected cyst, infection, infectious mononucleosis, injection site infection, localized infection, measles, oral candidiasis, orchitis, post procedural infection, postoperative wound infection, rhinitis, tonsillitis, urinary tract infection enterococcal

**INJURY, POISONING AND PROCEDURAL COMPLICATIONS:** arthropod bite, ligament sprain, post procedural complication, tibia fracture, wound, endotracheal intubation complication, fractured sacrum, gunshot wound, hand fracture, head injury, humerus fracture, joint injury, limb injury, meniscus lesion, mouth injury, muscle strain, nail injury, nerve injury, post procedural discomfort, post procedural hemorrhage, post-traumatic pain, procedural vomiting, rib fracture, spinal compression fracture, stress fracture, tooth injury, vena cava injury, wound dehiscence

**INVESTIGATIONS:** blood calcium decreased, blood pressure increased, breath sounds abnormal, neutrophil count decreased, blood iron decreased, blood lactate dehydrogenase

increased, c-reactive protein increased, cardiac murmur, computerized tomogram thorax abnormal, hemoglobin decreased, occult blood

**METABOLISM AND NUTRITION DISORDERS:** cell death, hyperlipidemia, iron deficiency

**MUSCULOSKELETAL AND CONNECTIVE TISSUE DISORDERS:** bursitis, joint range of motion decreased, joint warmth, tendonitis, wrist deformity, chondropathy, flank pain, intervertebral disc protrusion, joint effusion, limb discomfort, muscle contracture, muscle fatigue, osteoarthritis, osteonecrosis, peri-arthritis, plantar fasciitis, temporomandibular joint syndrome, tenosynovitis

**NEOPLASMS BENIGN, MALIGNANT AND UNSPECIFIED (INCL CYSTS AND POLYPS):** benign neoplasm of skin, skin papilloma, basal cell carcinoma, benign lung neoplasm, benign neoplasm, bone giant cell tumour, bone neoplasm, ganglioneuroma, lipoma, neoplasm progression, parathyroid tumour benign, sarcoma, spindle cell sarcoma, thyroid cancer, tumour hemorrhage

**NERVOUS SYSTEM DISORDERS:** presyncope, sinus headache, amnesia, carpal tunnel syndrome, central nervous system lesion, dysgeusia, hyperesthesia, intercostal neuralgia, lumbar radiculopathy, memory impairment, mental impairment, muscle contractions involuntary, nerve compression, nystagmus, peroneal nerve palsy, restless legs syndrome, tension headache, transient ischemic attack

**PSYCHIATRIC DISORDERS:** bruxism, libido decreased, mood altered, affect lability, depressed mood, emotional disorder, euphoric mood, mood swings, sleep disorder, stress, suicidal ideation

**RENAL AND URINARY DISORDERS:** pollakiuria, nocturia, polyuria, urine odour abnormal

**REPRODUCTIVE SYSTEM AND BREAST DISORDERS:** amenorrhea, breast cyst, breast discharge, breast pain, cervical dysplasia, dysmenorrhea, vaginal discharge, vaginal hemorrhage

**RESPIRATORY, THORACIC AND MEDIASTINAL DISORDERS:** paranasal sinus hypersecretion, pleuritic pain, rhinitis allergic, bronchial hyperreactivity, dyspnea exertional, hiccups, oropharyngeal blistering, pharyngeal disorder, productive cough, respiratory failure, respiratory tract congestion, sinus congestion, sneezing, upper-airway cough syndrome

**SKIN AND SUBCUTANEOUS TISSUE DISORDERS:** erythema, onychoclasia, scar pain, skin chapped, skin discoloration, skin hyperpigmentation, cold sweat, dermal cyst, dermatitis, dermatitis acneiform, dermatitis allergic, ecchymosis, keratosis pilaris, nail disorder, petechiae, pruritus generalized, rash erythematous, rash pruritic, seborrheic dermatitis, skin irritation, subcutaneous nodule

**VASCULAR DISORDERS:** flushing, lymphedema

\* The listing describes adverse events at initial integrated analysis (final analysis of Study 5 and interim analysis of Study 6) occurring in < 1% of patients in these studies.

## **8.4 Abnormal Laboratory Findings: Hematologic, Clinical Chemistry and Other Quantitative Data**

### **Clinical Trial Findings**

In clinical trials in cancer patients with bone metastases, a grade 3 decrease in serum calcium levels was experienced in 2.5% of patients treated with XGEVA and 1.2% of patients treated with zoledronic acid. A grade 4 decrease in serum calcium levels was experienced in 0.6% of patients treated with XGEVA and 0.2% of patients treated with zoledronic acid (see **7 WARNINGS AND PRECAUTIONS, Renal**). In studies, the development of hypocalcemia in patients with severe renal impairment (creatinine clearance less than 30 mL/min and/or receiving dialysis) was accompanied by an elevation in parathyroid hormone levels.

Severe hypophosphatemia (Grade 3) occurred in 15.4% of patients treated with XGEVA and 7.4% of patients treated with zoledronic acid.

In the clinical trials in patients with GCTB (final analysis of Study 5 and interim analysis of Study 6), the subject incidence of grade 2 corrected serum calcium decreases was 2.6%, while no grade 3 or grade 4 incidences were observed. In the same integrated analysis, Common Terminology Criteria for Adverse Events (CTCAE) grade 3 low phosphorus values were observed for 29 patients (9.5%).

In the final integrated analysis of both Studies 5 and 6, the subject incidence of grade 2, 3 and 4 corrected serum calcium decreases were 4.4%, 0.2% and 0.4%, respectively. CTCAE grade 3 low phosphorus values were observed for 108 patients (19.7%).

## **8.5 Post-Market Adverse Reactions**

In the post-marketing experience, the following have been reported in patients receiving XGEVA:

- Severe symptomatic hypocalcemia, including fatal cases
- Hypersensitivity, including anaphylactic reactions
- Musculoskeletal pain, including severe cases
- Lichenoid drug eruptions (eg, lichen planus-like reactions)
- Alopecia

## **9 DRUG INTERACTIONS**

### **9.1 Serious Drug Interactions**

No serious drug interactions have been identified.

### **9.2 Drug Interactions Overview**

No formal drug interaction studies have been conducted with XGEVA.

### **9.3 Drug-Behavioural Interactions**

Drug-behavioural interactions have not been established.

### **9.4 Drug-Drug Interactions**

Interactions with other drugs have not been established.

In clinical trials, XGEVA has been administered in combination with standard anti-cancer treatment and in patients previously receiving bisphosphonates. Apparent differences in the pharmacokinetics and pharmacodynamics of denosumab with concomitant chemotherapy and/or hormone therapy, or previous exposure to intravenous bisphosphonate were small in relation to inherent inter-subject variability within a patient population.

### **9.5 Drug-Food Interactions**

Interactions with food have not been established.

### **9.6 Drug-Herb Interactions**

Interactions with food herbal products have not been established.

### **9.7 Drug-Laboratory Test Interactions**

Interactions with laboratory tests have not been established.

## **10 CLINICAL PHARMACOLOGY**

### **10.1 Mechanism of Action**

XGEVA (denosumab) is a fully human IgG2 monoclonal antibody with high affinity and specificity for human RANK Ligand (RANKL). Binding of XGEVA to RANKL inhibits RANKL from activating its only receptor, RANK, on the surface of osteoclasts and their precursors. Increased osteoclast activity, stimulated by RANKL, is a key mediator of bone disease in metastatic tumours and multiple myeloma. Prevention of RANKL-RANK interaction inhibits osteoclast formation, function and survival, thereby decreasing bone resorption and interrupting cancer-induced bone destruction.

#### **Multiple Myeloma and Bone Metastasis from Solid Tumours**

XGEVA binds to RANK Ligand (RANKL), a transmembrane or soluble protein essential for the formation, function, and survival of osteoclasts, the cells responsible for bone resorption. XGEVA prevents RANKL from activating its receptor, RANK, on the surface of osteoclasts and their precursors. Increased osteoclast activity, stimulated by RANKL, is a key mediator of bone disease in metastatic tumours and multiple myeloma. Prevention of the RANKL/RANK interaction inhibits osteoclast formation, function, and survival, thereby decreasing bone resorption and interrupting cancer-induced bone destruction.

#### **Giant Cell Tumour of Bone**

GCTB are characterized by stromal cells expressing RANKL and osteoclast-like giant cells expressing RANK. In patients with GCTB, XGEVA binds to RANKL, significantly reducing or eliminating osteoclast-like giant cells. Consequently, osteolysis is reduced and proliferative tumour stroma is replaced with non-proliferative, differentiated, densely woven new bone.

#### **Hypercalcemia of Malignancy Refractory to Intravenous Bisphosphonates**

The primary etiology of both skeletal and humoral HCM is increased bone resorption, which leads to elevated calcium concentrations in the extracellular fluid. The increase in bone resorption is initiated by the release of signaling molecules such as PTHrP, prostaglandins, and cytokine by malignant and stromal cells. These molecules stimulate osteoblasts and other stromal cells to express RANK ligand (RANKL), which upon binding its receptor RANK



upregulates osteoclast recruitment and differentiation and thus bone resorption, with a resultant increase in calcium concentrations of the extracellular fluid and serum. XGEVA binds to RANKL preventing RANK/RANKL mediated osteoclast formation, function, and survival thereby lowering serum calcium levels.

### **Animal Pharmacology**

Denosumab has been shown to be a potent inhibitor of bone resorption in monkeys via inhibition of RANKL. Adolescent monkeys dosed with denosumab at 15 times (50 mg/kg dose) and 2.8 times (10 mg/kg dose) the area under the curve (AUC) exposure in adult humans dosed at 120 mg subcutaneously every 4 weeks had abnormal growth plates, considered to be consistent with the pharmacological activity of denosumab. Tissue distribution studies indicated that denosumab does not bind to tissues known for expression of other members of the TNF superfamily, including TNF-related apoptosis-inducing ligand (TRAIL).

Since the biological activity of denosumab in animals is specific to nonhuman primates, evaluation of genetically engineered (knockout) mice or use of other biological inhibitors of the RANK/RANKL pathway, such as OPG-Fc and RANK-Fc, were used to evaluate the pharmacodynamic properties of denosumab in rodent models. In mouse bone metastasis models of human prostate cancer, NSCLC, and estrogen receptor (ER) positive and negative breast cancer, OPG-Fc reduced osteolytic, osteoblastic, and osteolytic/osteoblastic lesions, delayed formation of de novo bone metastasis, and reduced skeletal tumour growth. When OPG-Fc was combined with hormonal therapy (tamoxifen) or chemotherapy (docetaxel) in these models, there was additive inhibition of skeletal tumour growth in breast, prostate or lung cancer respectively. In a mouse model of mammary tumour induction, RANK-Fc delayed tumour formation.

The role of osteoclast-mediated hypercalcemia was evaluated in 2 murine models of humoral hypercalcemia of malignancy through the use of osteoprotegerin (OPG), an endogenous decoy receptor that binds and neutralizes RANKL. In one model, mice were inoculated with syngeneic colon adenocarcinoma cells, and in the other mice were injected with high-dose parathyroid hormone-related protein (PTHrP) (0.5 mg/kg, SC, twice per day). In both models, a single injection of OPG caused more rapid reversal of established hypercalcemia and longer lasting suppression of hypercalcemia than high-dose bisphosphonates.

RANK/RANKL knockout mice exhibited absence of lymph node formation, as well as an absence of lactation due to inhibition of mammary gland maturation (lobulo-alveolar gland development during pregnancy). Neonatal RANK/RANKL knockout mice exhibited reduced bone growth and lack of tooth eruption. A corroborative study in 2-week-old rats given the RANKL inhibitor OPG-Fc also showed reduced bone growth, altered growth plates and impaired tooth eruption. These changes were partially reversible in this model when dosing with the RANKL inhibitors was discontinued (see **7 WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics**).

## 10.2 Pharmacodynamics

In a phase 2 study of patients with breast cancer and bone metastases who had not previously received intravenous (IV) bisphosphonate therapy, subcutaneous (SC) doses of XGEVA 120 mg every 4 weeks caused a rapid reduction in markers of bone resorption (uNTX/Cr and serum CTx) with a median reduction of 82% for uNTX/Cr within 1 week. Reductions in bone turnover markers were maintained, with median uNTX/Cr reductions of 74% to 82% from weeks 2 to 25 of continued 120 mg every 4 weeks dosing. In phase 3 clinical studies of patients with advanced cancer who had not previously received IV bisphosphonate therapy, median reductions of approximately 80% in uNTx/Cr from baseline after 3 months of treatment were observed across 2075 XGEVA-treated advanced cancer patients (breast, prostate, multiple myeloma or other solid tumours).

Similarly, in a phase 2 study of patients with solid tumours and bone metastases (including patients with multiple myeloma and bone disease) who were receiving IV bisphosphonate therapy, yet had uNTX/Cr levels > 50 nM/mM, SC dosing of XGEVA administered either every 4 weeks or every 12 weeks caused an approximate 80% reduction in uNTX/Cr from baseline after 3 and 6 months of treatment.

In a phase 3 study of patients with newly diagnosed multiple myeloma who received SC doses of XGEVA 120 mg every 4 weeks (Q4W), median reductions in uNTx/Cr of approximately 75% were observed by week 5. Reductions in bone turnover markers were maintained, with median reductions of 74% to 79% for uNTx/Cr from weeks 9 to 49 of continued 120 mg Q4W dosing.

In a phase 2 study of patients with GCTB who received SC doses of XGEVA 120 mg every 4 weeks (Q4W) with loading doses on days 8 and 15, median reductions in uNTx/Cr and serum CTx of approximately 80% were observed by week 9. Reductions in bone turnover markers were maintained, with median reductions of 56% to 77% for uNTx/Cr and 79% to 83% for serum CTx from weeks 5 to 25 of continued 120 mg Q4W dosing.

## 10.3 Pharmacokinetics

Following SC administration, bioavailability was 62% based on a population PK analysis. Relative AUC exposure ratios for SC vs. IV dosing were 78% and 75% for doses of 1.0 and 3.0 mg/kg in postmenopausal women. Denosumab displayed non-linear pharmacokinetics with dose over a wide dose range, but approximately dose-proportional increases in exposure for doses of 60 mg (or 1 mg/kg) and higher (for example, 3.8- to 4.0-fold increases in mean  $C_{max}$  and AUC values for a 3-fold increase in dose from 60 to 180 mg). In subjects with advanced cancer, who received multiple SC doses of 120 mg every 4 weeks an approximate 2.5-fold accumulation in serum denosumab  $AUC_{(0-\tau)}$  exposures was observed and steady-state was achieved on or after 6 doses. These results indicate that denosumab pharmacokinetics does not change with time or multiple dosing. In subjects with multiple myeloma who received 120 mg every 4 weeks, median trough levels varied by less than 8% between months 6 and 12. In subjects with GCTB who received 120 mg every 4 weeks with a loading dose on days 8 and 15, steady-state levels were achieved within the first month of treatment. Between weeks 9 and 49, median trough levels varied by less than 9%. At steady-state in these subjects, the mean serum trough concentration was 20.6 mcg/mL (range, 0.456 to 56.9 mcg/mL). In patients who discontinued 120 mg every 4 weeks dosing, the mean half-life was 28 days (range 14 to 55 days).

A population pharmacokinetic analysis was performed to evaluate the effects of demographic characteristics. This analysis suggested that there were no notable differences in various pharmacokinetics parameters (clearance, volume of distribution, absorption rate, bioavailability) with age (18 to 87 years), race, body weight (36 to 174 kg), or across patients with solid

tumours, multiple myeloma, and GCTB. Denosumab pharmacokinetics and pharmacodynamics were similar in men and women and in patients transitioning from IV bisphosphonate therapy. Denosumab pharmacokinetics and pharmacodynamics were not affected by the formation of binding antibodies to denosumab.

### **Special Populations and Conditions**

- **Pediatrics**

The pharmacokinetics of XGEVA has not been established in pediatric patients other than skeletally mature adolescents (aged 13-17 years) with GCTB.

In skeletally-mature adolescents (13 – 17 years of age; N = 10) with GCTB, the pharmacokinetics of denosumab were similar to those observed in adult subjects with GCTB (N = 15).

- **Geriatrics**

The pharmacokinetics of denosumab was not affected by age from 18 years to 87 years.

- **Sex**

The pharmacokinetics of denosumab was not different in men and women.

- **Ethnic Origin**

The pharmacokinetics of denosumab was not affected by race.

- **Hepatic Insufficiency**

No clinical studies have been conducted to evaluate the effect of hepatic impairment on the pharmacokinetics of denosumab.

- **Renal Insufficiency**

Two clinical trials were conducted in patients without cancer and with varying degrees of renal function. In one study, patients (N = 55) with varying degrees of renal function (ranging from normal through end-stage renal disease requiring dialysis) received a single 60 mg subcutaneous dose of denosumab. Hypocalcemia was observed in 8 subjects (15%), 1 (2%) of whom was symptomatic. Two patients (4%) each experienced an adverse event of hypocalcemia that was classified as serious. Both patients had severe chronic kidney disease (CKD) and were enrolled before the protocol required supplementation of calcium and vitamin D.

In a second study, patients (N = 32) with severe renal impairment (creatinine clearance less than 30 mL/minute and/or on dialysis) were given two 120 mg subcutaneous doses (Days 1 and 29) of denosumab. Two patients overall (1 in each group) had symptomatic hypocalcemia, based on clinical adverse events and concomitant symptoms. One patient in the severe CKD group had concomitant muscle spasms and 1 patient in the CKD on dialysis group had concomitant paresthesia. Hence, the overall incidence of clinically significant hypocalcemia (corrected serum calcium less than 1.75 mmol/L or symptomatic hypocalcemia) was 9.4%: 1 of 16 patients (6.3%) in the severe group and 2 of 16 patients (12.5%) in the CKD on dialysis group. Both events of symptomatic hypocalcemia were mild in severity.

Accompanying increases in parathyroid hormone have also been observed in patients receiving XGEVA with severe renal impairment or receiving dialysis.

In both studies in patients with varying degrees of renal function, including patients on dialysis, the degree of renal impairment had no effect on the pharmacokinetics and pharmacodynamics of denosumab. The risk of developing hypocalcemia increased with increasing degree of renal impairment, and in the absence of, or inadequate calcium supplementation.

Dose adjustment for renal impairment is not required and there is no additional need for more frequent renal monitoring due to XGEVA administration.

## **11 STORAGE, STABILITY AND DISPOSAL**

Store XGEVA in a refrigerator at 2°C to 8°C in the original carton. Do not freeze.

Prior to administration, XGEVA may be allowed to reach room temperature (up to 25°C) in the original container. Once removed from the refrigerator, XGEVA must not be exposed to temperatures above 25°C and must be used within 30 days. If not used within the 30 days, XGEVA should be discarded.

Do not use XGEVA after the expiry date printed on the label.

Protect XGEVA from direct light and heat.

Avoid vigorous shaking of XGEVA.

## **12 SPECIAL HANDLING INSTRUCTIONS**

Not applicable

## PART II: SCIENTIFIC INFORMATION

### 13 PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: denosumab

Molecular formula and molecular mass: 147 kDa

Structural formula: Denosumab is a fully human IgG2 monoclonal antibody heterotetramer consisting of two heavy chains of the gamma 2 subclass (447 amino acids per chain) and two light chains of the kappa subclass (215 amino acids per chain)

#### Product Characteristics:

Denosumab is produced in genetically engineered mammalian (Chinese hamster ovary) cells.

### 14 CLINICAL TRIALS

#### 14.1 Trial Design and Study Demographics

##### Bone Metastasis from Solid Tumours

**Table 6. Summary of Patient Demographics for Clinical Trials in Patients with Advanced Malignancies Involving Bone**

Study #	Trial design	Dosage, route of administration and duration*	Study subjects (n = number)	Mean age (Range)	Gender (Female:Male) %
Study 1	Phase 3, randomized, double-blind, active-controlled	XGEVA 120 mg SC and zoledronic acid placebo IV Q4W or zoledronic acid 4mg IV and denosumab placebo SC Q4W	2046 adults with advanced breast cancer and bone metastasis (XGEVA: 1026 Zoledronic acid: 1020)	57 (24, 91)	XGEVA (99.2:0.8) Zoledronic acid (99.1:0.9)
Study 2	Phase 3, randomized, double-blind, active-controlled	XGEVA 120 mg SC and zoledronic acid placebo IV Q4W or zoledronic acid 4mg IV and denosumab placebo SC Q4W	1776 adults with advanced cancers including solid tumours [excluding breast and prostate], multiple myeloma, and lymphoma (XGEVA: 886 Zoledronic acid: 890)	60 (18, 89)	XGEVA (33.6:66.4) Zoledronic acid (38.0:62.0)
Study 3	Phase 3, randomized, double-blind, active-controlled	XGEVA 120 mg SC and zoledronic acid placebo IV Q4W or zoledronic acid 4mg IV and denosumab placebo SC Q4W	1901 adult men with castrate-resistant prostate cancer and bone metastasis (XGEVA: 950 Zoledronic acid: 951)	71 (38, 93)	XGEVA (0:100) Zoledronic acid (0:100)

\* Studies were event-driven: the length of the primary double-blind treatment phase was determined by the anticipated date on which ~745 subjects experienced an initial on-study skeletal-related event.

The efficacy of XGEVA for the treatment of patients with advanced malignancies involving bone was demonstrated by three pivotal phase 3, international, randomized, double blind, active controlled studies compared with zoledronic acid: Study 1 in 2046 adults with advanced breast cancer and bone metastases; Study 2 in 1776 adults with other solid tumours [including non-small cell lung cancer (NSCLC), renal cell cancer, colorectal cancer, small cell lung cancer, bladder cancer, head and neck cancer, GI/genitourinary cancer and others, excluding breast cancer and prostate cancer] and bone metastases or multiple myeloma; and Study 3 in 1901 men with castrate-resistant prostate cancer and bone metastases.

Patients received either 120 mg XGEVA SC every 4 weeks or 4 mg zoledronic acid (dose-adjusted for reduced renal function) IV every 4 weeks. No dosage adjustments were necessary in patients receiving XGEVA. In accordance with the zoledronic acid prescribing information, patients with creatinine clearance < 30 mL/min were excluded. Daily supplements of  $\geq 500$  mg calcium and  $\geq 400$  IU of vitamin D were strongly recommended, unless hypercalcemia was present.

In each study, the primary outcome measure was to demonstrate non-inferiority of time to first on study skeletal-related event (SRE) as compared to zoledronic acid. The secondary outcome measures were superiority of time to first on-study SRE and superiority of time to first and subsequent SRE; testing for the secondary outcome measures occurred if the primary outcome measure was statistically significant. An SRE is defined as any of the following: pathologic fracture, radiation therapy to bone, surgery to bone or spinal cord compression.

### **Multiple Myeloma**

In Study 4, XGEVA was evaluated in an international, randomized (1:1), double-blind, active-controlled study comparing XGEVA with zoledronic acid in patients with newly diagnosed multiple myeloma.

In this study, 1718 multiple myeloma patients with at least 1 bone lesion were randomized to receive 120 mg XGEVA subcutaneously every 4 weeks or 4 mg zoledronic acid intravenously (IV) every 4 weeks (dose adjusted for renal impairment and patients with creatinine clearance less than 30 mL/min were excluded based on zoledronic acid prescribing information). Randomization was stratified by intent to undergo autologous peripheral blood stem cell (PBSC) transplantation (yes or no), the anti-myeloma agent being utilized/planned to be utilized in first-line therapy [novel therapy-based or non-novel therapy-based (novel therapies include bortezomib, lenalidomide, or thalidomide)], stage at diagnosis (International Staging System I or II or III), previous SRE (yes or no), and region (Japan or other countries). The primary outcome measure was demonstration of non-inferiority of time to first skeletal-related event (SRE) as compared to zoledronic acid. Secondary outcome measures included superiority of time to first SRE, superiority of time to first and subsequent SRE, and overall survival. An SRE was defined as any of the following: pathologic fracture (vertebral or non-vertebral), radiation therapy to bone (including the use of radioisotopes), surgery to bone, or spinal cord compression.

Across both study arms, 54.5% of patients intended to undergo autologous PBSC transplantation, 95.8% patients utilized/planned to utilize a novel anti-myeloma agent in first-line therapy, and 60.7% of patients had a previous SRE. The number of patients across both study arms with ISS stage I, stage II, and stage III at diagnosis were 32.4%, 38.2%, and 29.3%, respectively.

Median age was 63 years (range: 29 – 91), 82% of patients were White, 12% Asian, 4% Black, and 46% of patients were women. The median number of doses administered was 16 for XGEVA and 15 for zoledronic acid.

## Giant Cell Tumour of Bone

**Table 7. Summary of Patient Demographics for Clinical Trials in Patients with Giant Cell Tumour of Bone (GCTB)**

Study #	Trial design	Dosage, route of administration	Study subjects (n = number)	Mean age (Range) <sup>a</sup>	Gender (Female:Male) % <sup>a</sup>
Study 5	Phase 2, open-label, multicenter	XGEVA 120 mg SC Q4W with a loading dose on study days 8 and 15 of the first month of therapy	Adult subjects with GCTB (n = 37)	34 (19, 63)	(54:46)
Study 6 (interim analysis)	Phase 2, open-label, multicenter	XGEVA 120 mg SC Q4W with a loading dose on study days 8 and 15 of the first month of therapy	Adult subjects with GCTB (n = 272) And Skeletally Mature Adolescents with GCTB (n = 10)	36 (13, 83)	(58:42)

<sup>a</sup> Pooled data

The safety and efficacy of XGEVA was studied in two phase 2 open-label, single arm trials (Studies 5 and 6) that enrolled patients with GCTB that was either unresectable or for which surgery was associated with severe morbidity.

Study 5 enrolled 37 adult patients with histologically confirmed unresectable or recurrent GCTB and the main outcome measure of the trial was response rate based on histological or radiographic evidence.

At the interim analysis, Study 6 enrolled 282 adults, and 10 skeletally mature adolescents (aged 13-17 years) with GCTB. By the end of the study, Study 6 had enrolled 507 adults and 28 skeletally mature adolescents (aged 13-17 years) with GCTB. The main outcome measure was to evaluate the safety profile of XGEVA. Efficacy was assessed by evaluation of time to disease progression in subjects with unresectable GCTB and by evaluation of the proportion of subjects who do not require surgery in the subjects with resectable GCTB.

A retrospective independent review of radiographic imaging data from 190 of 305 patients enrolled in Studies 5 and 6 was performed. Patients were evaluated using modified Response Evaluation Criteria in Solid Tumours (RECIST 1.1) to evaluate tumour burden based on computed tomography (CT)/magnetic resonance imaging (MRI).

### Hypercalcemia of Malignancy Refractory to Intravenous Bisphosphonate

The safety and efficacy of XGEVA was studied in a phase 2 open-label, single-arm trial (Study 7) that enrolled 33 patients with hypercalcemia of malignancy (with or without bone metastases) refractory to treatment with intravenous bisphosphonate. In this study, refractory hypercalcemia of malignancy was defined as an albumin-corrected serum calcium (CSC) of >12.5 mg/dL (3.1 mmol/L) despite treatment with intravenous bisphosphonate in the last 7-30 days. Patients receiving dialysis for renal failure or who had treatment with thiazides, calcitonin, mithramycin, or gallium nitrate within their window of expected therapeutic effect prior to the date of screening corrected serum calcium (CSC) were excluded. Twenty-six (79%) patients had advanced solid tumours and 7 (21%) patients had advanced hematologic malignancies. Twenty-five patients (76%) had poor performance status (Eastern Cooperative Oncology Group

[ECOG]  $\geq 2$ ) at baseline. Metastatic disease was present in 30 (91%) patients and metastatic bone disease in 13 (39%) patients at baseline. Three (9%) patients had non-metastatic disease, 2 with myeloma and 1 with non-Hodgkin's lymphoma.

At the time of enrollment, the median serum calcium level was 13.7 mg/dL (3.42 mmol/L). During the study, serum calcium was collected every few days in the first month, weekly during the second month, and monthly thereafter.

The primary outcome measure was the proportion of patients achieving a response, defined as CSC  $\leq 11.5$  mg/dL (2.9 mmol/L), within 10 days after XGEVA administration. Additional secondary outcomes were assessed and are presented in Table 11.

Patients received XGEVA subcutaneously every 4 weeks with additional 120 mg doses on days 8 and 15 of the first month of therapy.

## **14.2 Study Results**

### **Bone Metastasis from Solid Tumours**

XGEVA reduced the risk of developing (delayed time to) first SRE and multiple (first and subsequent) SREs in patients with advanced malignancies involving bone. Efficacy results are provided in Table 8.



**Table 8. Efficacy Results for XGEVA Compared to Zoledronic Acid in Patients with Advanced Malignancies Involving Bone**

	Study 1 Advanced Breast Cancer		Study 2 Advanced Cancer (Other Solid Tumours and Multiple Myeloma)		Study 3 Advanced Prostate Cancer	
	XGEVA	Zoledronic Acid	XGEVA	Zoledronic Acid	XGEVA	Zoledronic Acid
N	1026	1020	886	890	950	951
<b>First On-Study Skeletal Related Event (SRE)</b>						
Number and Proportion of Subjects with SREs (%)	315 (30.7)	372 (36.5)	278 (31.4)	323 (36.3)	341 (35.9)	386 (40.6)
Components of First SRE						
Radiation to Bone	82 (8.0)	119 (11.7)	119 (13.4)	144 (16.2)	177 (18.6)	203 (21.3)
Pathological Fracture	212 (20.7)	238 (23.3)	122 (13.8)	139 (15.6)	137 (14.4)	143 (15.0)
Surgery to Bone	12 (1.2)	8 (0.8)	13 (1.5)	19 (2.1)	1 (0.1)	4 (0.4)
Spinal Cord Compression	9 (0.9)	7 (0.7)	24 (2.7)	21 (2.4)	26 (2.7)	36 (3.8)
Median Time (months)	NR	26.4	20.5	16.3	20.7	17.1
Hazard Ratio (95% CI)	0.82 (0.71, 0.95)		0.84 (0.71, 0.98)		0.82 (0.71, 0.95)	
Non-inferiority P-value	<0.0001		0.0007		0.0002	
Superiority P-value †	0.0101		0.0619		0.0085	
<b>First and Subsequent SRE *</b>						
Mean Number/Patient	0.46	0.60	0.44	0.49	0.52	0.61
Rate Ratio (95% CI)	0.77 (0.66, 0.89)		0.90 (0.77, 1.04)		0.82 (0.71, 0.94)	
Superiority P-value †	0.0012		0.1447		0.0085	

NR = not reached

CI = confidence interval

Superiority testing performed only after denosumab demonstrated to be non-inferior to zoledronic acid within trial.

\*Accounts for all skeletal events over time; only events occurring  $\geq$  21 days after the previous event are counted.

†P-values, adjusted for multiplicity, are presented for Studies 1, 2 and 3

Overall survival and disease progression in all three studies were comparable in patients with advanced cancer between XGEVA and zoledronic acid treatment groups (see Table 9). In Study 2, mortality was higher with XGEVA in a subgroup analysis of patients with multiple myeloma [hazard ratio (95% CI) of 2.26 (1.13, 4.50); n = 180].

**Table 9. Summary of Exploratory Tumour Outcomes**

Endpoint	XGEVA vs Zoledronic acid Hazard Ratio					
	Study 1		Study 2		Study 3	
	Pt Est	95% CI*	Pt Est	95% CI*	Pt Est	95% CI*
Overall survival	0.95	0.81, 1.11	0.95	0.83, 1.08	1.03	0.91, 1.17
Time to disease progression excluding death	1.00	0.89, 1.11	1.00	0.89, 1.12	1.06	0.95, 1.18

Pt Est = point estimate

CI = confidence interval

\*Not adjusted for multiplicity

**Multiple Myeloma**

In patients with newly diagnosed multiple myeloma, XGEVA was non-inferior to zoledronic acid in delaying the time to first SRE following randomization (Table 10).

**Table 10. Efficacy Results for XGEVA Compared to Zoledronic Acid in Patients with Newly Diagnosed Multiple Myeloma**

	XGEVA (N = 859)	Zoledronic Acid (N = 859)
<b>First SRE</b>		
Number of Patients who had SREs (%)	376 (43.8)	383 (44.6)
Median Time to SRE (months) (95% CI)	22.83 (14.72, NE)	23.98 (16.56, 33.31)
Hazard Ratio <sup>a</sup> (95% CI)	0.98 (0.85, 1.14)	
Non-inferiority p-value <sup>a</sup>	0.010	
<b>Components of First SRE</b>		
Radiation to Bone	47 (5.5)	62 (7.2)
Pathological Fracture	342 (39.8)	338 (39.3)
Surgery to Bone	37 (4.3)	48 (5.6)
Spinal Cord Compression	6 (0.7)	4 (0.5)

<sup>a</sup>Based on a Cox proportional hazards model stratified by randomization stratification factors

The hazard ratio between XGEVA and zoledronic acid treatment groups and 95% CI for overall survival (OS) was 0.90 (0.70, 1.16).


**Giant Cell Tumour of Bone**

Overall, in the retrospective interim analysis, an objective response by RECIST 1.1 was observed in 47 of 187 (25.1%) evaluable patients (95% CI: 19.1, 32.0), including 2 of 6 (33.3%) evaluable adolescent patients. All responses were partial responses. The median time to response was 3 months (range: 1.5 to 20.9 months). The median duration of response was not estimable as only three patients experienced disease progression following an objective response. The median follow-up duration for evaluable patients was 13 months (range: 2 to 49 months).

## Hypercalcemia of Malignancy Refractory to Intravenous Bisphosphonate

Based on Kaplan-Meier estimates, 59% of patients met the primary end point by day 10 and 73.9% by day 57, responding to XGEVA treatment with CSC levels  $\leq 11.5$  mg/dL.

**Figure 1. Corrected serum calcium by visit (median and interquartile range)**



N = Number of patients who received at least 1 dose of denosumab; n = Number of patients who had no missing data at baseline and the timepoint of interest

**Table 11. Efficacy in Patients with Hypercalcemia of Malignancy Refractory to Bisphosphonate Therapy**

	N = 33	Kaplan-Meier Estimates by Visit (95% CI <sup>a</sup> )
<b>All Responders (CSC <math>\leq 11.5</math> mg/dL) by Day 10</b>	21	59.0% (41.5%, 74.5%)
<b>All Responders by Day 57</b>	23	73.9% (56.7%, 86.2%)
<b>Complete Responders (CSC <math>\leq 10.8</math> mg/dL) by Day 10</b>	12	34.3% (19.3%, 52.7%)
<b>All Complete Responders by Day 57</b>	21	75.2% (56.3%, 88.4%)

N = Number of subjects who received  $\geq 1$  investigational product and have a screening serum calcium corrected by albumin (from local lab)  $> 12.5$  mg/dL (3.1 mmol/L).

CSC = corrected serum calcium

<sup>a</sup> Confidence interval is calculated using bootstrap method

Median time to response (CSC  $\leq$  11.5 mg/dL) was 9 days (95% CI: 8, 19), and the median duration of response was 104 days (95% CI: 7, not estimable). Median time to complete response (CSC  $\leq$  10.8 mg/dL) was 23 days (95% CI: 9, 36), and the median duration of complete response was 34 days (95% CI: 1, 134).

### 14.3 Comparative Bioavailability Studies

Not Applicable.

### 14.4 Immunogenicity

Denosumab is a human monoclonal antibody. As with all therapeutic proteins, there is potential for immunogenicity. In clinical studies, neutralizing antibodies have not been observed for XGEVA in advanced cancer patients or GCTB patients. Using an electro chemiluminescent bridging immunoassay, less than 1% of patients treated with XGEVA for up to 3 years tested positive for binding antibodies (including pre-existing, transient, and developing antibodies). None of the patients tested positive for neutralizing antibodies as assessed using a chemiluminescent cell-based *in vitro* biological assay. There was no evidence of altered pharmacokinetic profile, toxicity profile, or clinical response associated with binding antibody development.

The incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of a positive antibody (including neutralizing antibody) test result may be influenced by several factors, including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of antibodies to denosumab with the incidence of antibodies to other products may be misleading.

## 15 MICROBIOLOGY

No microbiological information is required for this drug product.

## 16 NON-CLINICAL TOXICOLOGY

### Carcinogenicity

Since denosumab is highly species-specific and is not active in rodents, traditional rodent cancer bioassays could not be performed. RANKL inhibition (the target of denosumab) has been studied in a wide range of short-term animal models of cancer and shown no carcinogenic potential. Additionally, RANKL inhibition has shown no evidence of immunosuppression in a wide range of animal models.

### Genotoxicity

The genotoxic potential of denosumab has not been evaluated. Denosumab is a recombinant protein made up entirely of naturally-occurring amino acids and contains no inorganic or synthetic organic linkers or other non-protein portions. Therefore, it is unlikely that denosumab or any of its derived fragments would react with DNA or other chromosomal material.

## **Reproductive and Developmental Toxicology**

Denosumab had no effect on female fertility or male reproductive organs in monkeys at exposures that were 9.5- to 16-fold higher than the human exposure for 120 mg SC administered once every 4 weeks.

**Table 12. Summary of Preclinical Toxicity and Reproductive Studies with Denosumab**

Type of Study	Species and strain	Number per sex per group	Route of Administration	Dose (mg/kg) and dosing regimen	Study Duration	Treatment related findings	NOAEL (mg/kg)
Repeated-dose Toxicity	Cynomolgus monkey	6	Subcutaneous or Intravenous	Once weekly: 0, 0.1, 1.0, & 10.0 (SC); 10.0 (IV)	1-month dosing with 3 months recovery	<p>Consistent with the pharmacological action of denosumab, there were rapid and marked decreases in circulating markers of bone turnover at all doses. Correlating with these changes, there was increased bone mineral density in males dosed at 1 and 10 mg/kg. With the exception of bone mineral density which tended to be maintained, these changes were recovered or recovering following 3 treatment free months.</p> <p>There were no treatment related effects on organ weights or histopathology findings.</p>	10 (SC and IV)

**Table 12. Summary of Preclinical Toxicity and Reproductive Studies with Denosumab**

Type of Study	Species and strain	Number per sex per group	Route of Administration	Dose (mg/kg) and dosing regimen	Study Duration	Treatment related findings	NOAEL (mg/kg)
	Cynomolgus monkey	8	Subcutaneous	Once monthly: 0, 1, 10, 50	6 and 12 months with 3 months recovery	Consistent with the pharmacological action of denosumab, there were rapid and marked decreases in circulating markers of bone turnover at 10 and 50 mg/kg. Correlating to these changes, there was increased bone mineral density, bone mineral content, cortical area and thickness, and bone strength parameters in males dosed at 50 mg/kg, and females dosed at 10 and 50 mg/kg. In addition, there was enlargement of the growth plates, decreased osteoblasts and osteoclasts, and decreased chondroclasis at 10 and 50 mg/kg. These changes were recovered or recovering following 3 treatment free months. There were no treatment related changes in ophthalmoscopy, cardiovascular physiology, sperm motility and morphology, circulating immunoglobulins and lymphocyte subsets, or organ weights.	50
Female Fertility	Cynomolgus monkey	6 Females	Subcutaneous	Once weekly: 0, 2.5, 5, 12.5	Over 2 menstrual cycles before mating and for 4 weeks after mating	No treatment related effects on cyclicity, circulating reproductive hormones, mating success.	12.5

**Table 12. Summary of Preclinical Toxicity and Reproductive Studies with Denosumab**

Type of Study	Species and strain	Number per sex per group	Route of Administration	Dose (mg/kg) and dosing regimen	Study Duration	Treatment related findings	NOAEL (mg/kg)
Embryo-fetal Development	Cynomolgus monkey	16 Females	Subcutaneous	Once weekly: 0, 2.5, 5, 12.5	Gestation days 20-50	No treatment related effects on mother or embryonic development were observed. Peripheral lymph nodes were not evaluated.	12.5
Enhanced pre- and post-natal development	Cynomolgus monkey	29 Females	Subcutaneous	Once monthly: 0, 50	Gestation days 20 - 22 to birth	There were increased fetal losses during gestation, increased stillbirths and post-natal mortality (see Table 13). Treatment-related findings in the offspring included decreased body weight gain and decreased neonatal growth; skeletal abnormalities resulting from impaired bone resorption during rapid growth, including bones at the base of the skull resulting in altered cranial shape and exophthalmos, reduced bone strength and treatment-related bone fractures; reduced hematopoiesis; decreased serum levels of bone resorption and bone formation biomarkers; tooth malalignment and dental dysplasia (in the absence of adverse effects on tooth eruption); infections; and absence of peripheral lymph nodes.	A NOAEL was not identified.



**Table 12. Summary of Preclinical Toxicity and Reproductive Studies with Denosumab**

Type of Study	Species and strain	Number per sex per group	Route of Administration	Dose (mg/kg) and dosing regimen	Study Duration	Treatment related findings	NOAEL (mg/kg)
						<p>Following a recovery period from birth out to 6 months of age, findings still observed were mildly reduced bone length (femoral, vertebral, jaw); reduced cortical thickness with associated reduced strength; extramedullary hematopoiesis; dental dysplasia; and the absence of decreased size of some lymph nodes. One infant had minimal to moderate mineralization in multiple tissues. The initially lower growth rates returned to, but never exceeded the growth rate in the control group, and hence, the infants exposed to denosumab remained smaller than control infants, as measured by body weight and morphometric measurements. For the denosumab-treated maternal animals, there was a decrease in serum levels of bone resorption and formation biomarkers, and serum alkaline phosphatase levels; recovery was evident by the end of the treatment-free period. Maternal mammary gland development was normal. At birth out to 1 month of age, infants had measurable blood levels of denosumab (22-621% of the maternal levels). Only one infant had measurable concentrations of denosumab on BD91, and no infants had measurable concentrations on BD180. Generally, the effects observed in mothers and infants were consistent with the pharmacological action of denosumab.</p>	

**Table 12. Summary of Preclinical Toxicity and Reproductive Studies with Denosumab**

Type of Study	Species and strain	Number per sex per group	Route of Administration	Dose (mg/kg) and dosing regimen	Study Duration	Treatment related findings	NOAEL (mg/kg)
Safety Pharmacology	Cynomolgus monkey	3 Males	Subcutaneous	Single dose: 0, 0.3, 3, 30	7 Days	No treatment related effects on heart rate, blood pressure, electrical activity of the heart, or respiratory rate were observed.	30
	Sprague Dawley weanling rats	71 males and 67 females	Subcutaneous	Rat OPG-Fc: 1, 10 mg/kg/week Murine RANK-Fc: 10 mg/kg/week	6 weeks	Increased bone volume, density and strength. Increased cancellous bone with reduced osteoclast number. Reduced long bone growth with altered growth plate morphology and increased thickness. Impaired tooth eruption and tooth root formation.	N/A
		10 males and 3-10 females	Subcutaneous	Rat OPG-Fc: 3, 10 mg/kg/week	6 weeks	Changes seen at the 10 mg/kg/week were similar to those in the previous study. Effects were less at the 3 mg/kg/week.	N/A
		10-11 males and 9-10 females	Subcutaneous	Rat OPG-Fc: 1, 3, 10 mg/kg/week	6 weeks with 10 weeks recovery	Effects were partially reversible when OPG-Fc was discontinued	N/A
Other Studies – Tissue Cross-reactivity	Cynomolgus monkey, rat, rabbit	N/A	<i>In Vitro</i>	5 or 25 mcg/mL	N/A	Staining of lymphoid tissue in rabbit and cynomolgus monkey and staining of chondrocytes in rat were observed.	N/A
	Cynomolgus monkey, human	N/A	<i>In Vitro</i>	1 or 10 mcg/mL	N/A	Staining of lymphoid tissue in monkey, but no staining in human tissue was observed.	N/A
	Human	N/A	<i>In Vitro</i>	1 or 10 mcg/mL	N/A	Staining of lymphoid tissue was observed.	N/A

N/A = not applicable; NOAEL = no observed adverse effect level

**Table 13. Total Fetal Losses<sup>c</sup>, all Groups**

Dose (mg/kg)	Total No. Pregnant Females; Infants Born (M/F)	Gestation Day (GD) of Fetal Loss	% Fetal Loss by Dose Level			
			Full Gestation	First Trimester (GD20 to GD50)	Third Trimester Total (≥GD100)	Third Trimester Stillbirths (≥GD140)
0	29; 22 (13/9)	GDs 32, 32, 33, 104, 152, 157, 170	24.1% (7/29)	10.3% (3/29)	13.8% (4/29)	10.3% (3/29)
50	29; 16 (7/9)	GDs 31, 32, 33, 33, 46, 88 <sup>a</sup> , 132, 151, 156 <sup>a</sup> , 157, 158, 160, 168	40.7% (11/27) 44.8%** (13/29)	17.2% (5/29)	22.2% (6/27) 24.1%** (7/29)	18.5% (5/27) 20.7%** (6/29)
Historical Control Data <sup>b</sup>			24.8% (33/133)	6.8% (9/133)	15.8% (21/133)	9.0% (12/133)
Range			(6.7 to 38.9%)	(0 to 11.8%)	(0 to 28.6%)	(0 to 16.7%)

<sup>a</sup> Two adult females were excluded from fetal loss calculations except for first trimester because each had an anti-drug antibody (ADA) response beginning at GD76 with subsequent decrease in pharmacologic effect (bone biomarkers) prior to fetal loss; results indicated by a double asterisk (\*\*) include these ADA-positive adult females

<sup>b</sup> Based on 8 enhanced PPND studies conducted at the Testing Facility from 2008 to 2010

<sup>c</sup> Fetal losses occurring prior to GD140 were considered abortions; those occurring on or after GD140 were considered stillbirths

## PATIENT MEDICATION INFORMATION

### READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

<sup>Pr</sup>**XGEVA**<sup>®</sup> (*ex-jee-va*)

#### **denosumab injection**

Read this carefully before you start taking **XGEVA** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **XGEVA**.

#### **Serious Warnings and Precautions**

- Osteonecrosis of the jaw (sore in mouth involving gums or jaw bones)

#### **What is XGEVA used for?**

- XGEVA is used for reducing the risk of developing cancer-related complications like broken bones and/or bone pain that need surgery or radiation.
- XGEVA is used for reducing the risk of developing cancer-related complications in patients with multiple myeloma. Multiple myeloma is a cancer of plasma cells (a type of white blood cell).
- XGEVA is used to treat giant cell tumour of bone, which cannot be treated by surgery or where surgery is not the best option in adults and adolescents (aged 13-17 years) whose bones have stopped growing.
- XGEVA is used to reduce high levels of calcium in the blood in cancer patients (hypercalcemia of malignancy) after other drugs called bisphosphonates did not work.

#### **How does XGEVA work?**

XGEVA works differently than other medications used to treat cancer patients whose disease has spread to their bones. It works as a RANK Ligand (RANKL) inhibitor. RANKL is a protein that promotes the breakdown of bone. XGEVA blocks RANKL to stop the breakdown of bone. This action strengthens your bones by increasing bone mass and lowers the chance of the cancer causing problems with your bones, such as fractures or severe pain requiring radiation treatment.

XGEVA reduces the amount of calcium in the blood by reducing the breakdown of bones. In patients with hypercalcemia of malignancy, the breakdown of bones can cause too much calcium in the blood.

#### **What are the ingredients in XGEVA?**

Medicinal ingredients: denosumab

Non-medicinal ingredients: acetate, polysorbate 20, sodium hydroxide, sorbitol and water for injection.

**XGEVA comes in the following dosage forms:**

XGEVA is a liquid for injection, with enough liquid in it for one shot. Each vial delivers 120 mg of denosumab. XGEVA is supplied in a carton containing 1 vial.

**Do not use XGEVA if:**

- you are allergic to denosumab or any other ingredient of XGEVA. Allergic reactions (eg, rash, hives, or in rare cases, swelling of the face, lips, tongue, throat, or trouble breathing) have been reported.
- you have hypocalcemia (low calcium levels in the blood), until your doctor corrects this condition.

**What important information do I need to know about taking XGEVA?**

XGEVA contains the same medicine as another drug called PROLIA, but at a different dose. If you are being treated with XGEVA, you should not be taking PROLIA or vice versa.

**To help avoid side effects and ensure proper use, talk to your healthcare professional before you take XGEVA. Talk about any health conditions or problems you may have, including:**

**Hypocalcemia (low calcium levels in the blood)**

XGEVA may lower levels of calcium in your blood. In the post-marketing setting, cases of low blood calcium with severe symptoms, including death, have been reported. If you have low blood calcium before you start receiving XGEVA, it may get worse during treatment. Your low blood calcium must be treated before you receive XGEVA. Most people with low calcium levels do not have symptoms, but some people may have symptoms. When the calcium levels in your blood go down, your body tries to correct the calcium levels by increasing the amount of a hormone made by your parathyroid glands (parathyroid hormone) in your blood. Call your doctor right away if you have symptoms of low blood calcium such as:

- Spasms, twitches, or cramps in your muscles.
- Numbness or tingling in fingers, toes or around the mouth.

Conditions which increase the risk of low blood calcium:

- If you cannot take daily calcium and/or vitamin D.
- If you have severe kidney disease or are on dialysis.

Your doctor will tell you to take calcium and vitamin D to help prevent low calcium levels in your blood while you take XGEVA, unless your blood calcium is high. Take calcium and vitamin D as your doctor tells you to.

**Osteonecrosis of the Jaw (sore in mouth involving gums or jaw bones)**

Severe jaw bone problems may happen when you take XGEVA. Your doctor should examine your mouth before you start XGEVA. Your doctor may tell you to see your dentist before you start XGEVA. It is important for you to practice good mouth care such as brushing and flossing your teeth regularly during treatment with XGEVA.

Tell your doctor immediately about any dental symptoms, including pain or unusual feeling in your teeth or gums, or any dental infections. If possible, you should not undergo tooth

extraction or other dental procedures (excluding regular dental cleaning) while you are receiving treatment with XGEVA without talking to your doctor first.

If you do need dental work, tell your dentist that you are receiving XGEVA and tell your doctor that you are having dental work done.

### **Unusual Thigh Bone Fractures**

Unusual fracture in the thigh bone may occur with some medicines, including XGEVA. Contact your doctor if you experience new or unusual pain in your hip, groin, or thigh.

### **High Calcium Levels in the Blood after Stopping Treatment with XGEVA**

Some patients with giant cell tumour of the bone and some who are still growing during treatment with XGEVA, have developed high calcium levels in the blood weeks to months after stopping treatment.

Your doctor will monitor you for signs and symptoms of high levels of calcium, after you stop receiving XGEVA.

### **Risk of Broken Bones in the Spine After Stopping Treatment with XGEVA**

Do not stop taking XGEVA without first talking with your doctor. After treatment with XGEVA is stopped, there may be an increased risk of having broken bones in your spine especially in people who have had a fracture or who have had osteoporosis (a condition in which bones become thin and fragile).

### **Skin Infections**

Tell your doctor promptly if you develop a swollen, red area on your skin that feels hot and tender with symptoms of fever (cellulitis) while taking XGEVA.

### **Pregnancy or Breast-Feeding**

XGEVA is not recommended for use in women who are pregnant or plan to become pregnant and nursing mothers. XGEVA may interfere with normal bone and tooth development in fetuses and nursing babies, and may interfere with breastfeeding.

XGEVA is not intended for use in pregnant women. You should not be given XGEVA if you are pregnant. A highly effective method of birth control should be used when taking XGEVA, or for at least 5 months after the last dose of XGEVA.

It is not known whether XGEVA is excreted into human milk.

### **Use in Children**

XGEVA is not recommended for anyone under 18 years of age except for adolescents with giant cell tumour of bone whose bones have stopped growing. The use of XGEVA has not been studied in children and adolescents with other cancers that have spread to bone.

**Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.**

### **The following may interact with XGEVA:**

Before starting XGEVA, tell your doctor about all the medicines you take, including prescription and non-prescription drugs, vitamins and herbal supplements.

Interactions between XGEVA and other drugs have not been studied.

### **How to take XGEVA:**

XGEVA is administered as a single injection under the skin (subcutaneous) once every four weeks. You should not inject XGEVA into the muscle (intramuscular), into your veins (intravenous) or between the layers of the skin (intra-dermal). The injection can be in your upper arm, upper thigh, or abdomen. The injection should be administered under the supervision of your doctor who is familiar with this drug. You may be able to give yourself XGEVA injections only if you have been trained in giving the injection and your doctor thinks you are capable of doing it correctly and if your doctor follows up with you as necessary.

Before injection, remove the vial from the refrigerator and allow it to reach room temperature (up to 25°C) in the original container. This will make the injection more comfortable. Do not shake. See instructions for injection.

Keep all medicines, including XGEVA, away from children.

Do not share XGEVA product with others, even if they have a similar disease.

### **INSTRUCTIONS FOR INJECTION**

**IMPORTANT:** TO HELP AVOID CONTAMINATION AND POSSIBLE INFECTION DUE TO INJECTION, PLEASE READ AND FOLLOW THESE INSTRUCTIONS CAREFULLY.

#### **How to prepare for XGEVA injection**

XGEVA is available as a liquid in vials. When you receive your XGEVA, always check to see that:

- The name XGEVA appears on the package and vial label.
- The expiration date on the vial label has not passed. **Do not use a vial after the date on the label.**
- The XGEVA liquid in the vial is clear, colourless to slightly yellow.

Only use disposable syringes and needles. Use the syringes only once and dispose of them as instructed by your doctor or nurse.

#### **Setting up for an injection**

1. Find a clean flat working surface, such as a table.
2. Remove the vial of XGEVA from the refrigerator. Allow XGEVA to reach room temperature (this takes about 15 to 30 minutes). Vials should be used only once. **DO NOT SHAKE THE VIAL.** Shaking may damage the XGEVA. If the vial has been shaken vigorously, the solution may appear foamy and it should not be used.
3. Assemble the supplies you will need for an injection:
  - XGEVA vial and sterile disposable syringe and a 27-gauge needle.
  - Two alcohol swabs and one cotton ball or gauze pad.
  - Puncture-proof disposal container.
4. Clean your work surface thoroughly and wash your hands with soap and warm water.

## Selecting and preparing the injection site


1. Choose an injection site. The recommended injection sites for XGEVA are:
  - The outer area of your upper arms.
  - The abdomen, except for the two-inch (5 cm) area around your navel.
  - The top of your thighs.

## How to prepare the dose of XGEVA in vials

1. Take the cap off the vial. Clean the stopper with an alcohol swab.
2. Check the package containing the syringe. If the package has been opened or damaged, do not use that syringe. Dispose of that syringe in the puncture-proof disposal container. If the syringe package is undamaged, open the package and remove the syringe.
3. Keep the vial on your flat working surface and insert the needle straight down through the rubber stopper. Do not put the needle through the rubber stopper more than once.
4. Push the plunger of the syringe down and inject the air from the syringe into the vial of XGEVA. Keeping the needle inside the vial, turn the vial upside down. Make sure that the tip of the needle is in the XGEVA liquid.
5. Keeping the vial upside down, slowly pull back on the plunger to fill the syringe with XGEVA liquid. Withdraw the entire content of the vial.
6. Keeping the needle in the vial, turn the syringe needle up and check for air bubbles in the syringe. If there are air bubbles, gently tap the syringe with your fingers until the air bubbles rise to the top of the syringe. Then slowly push the plunger up to force the air bubbles out of the syringe.
7. Remove the syringe from the vial but **do not lay it down** or let the needle touch anything.

## Injecting the dose of XGEVA

1. Hold the syringe in the hand you will use to inject XGEVA. With the other hand, clean the injection site with an alcohol swab. Use a circular motion from the inside to the outside of the injection site.
2. Pinch a fold of skin at the cleaned injection site.
3. Holding the syringe like a pencil, use a quick “dart-like” motion to insert the needle either straight up and down (90-degree angle) or at a slight angle (45 degrees) into the skin.





4. After the needle is inserted, let go of the skin. Inject the prescribed dose subcutaneously as directed by your doctor, nurse or pharmacist.
5. When the syringe is empty, pull the needle out of the skin and place a cotton ball or gauze over the injection site and press for several seconds.
6. Use a syringe, needle and vial only once. **DO NOT** put the needle cover (the cap) back on the needle. Discard the vial with any remaining XGEVA liquid.

### **Disposal of syringes, needles and vials**

You should always follow the instructions given by your doctor, nurse, or pharmacist on how to properly dispose of containers with used syringes, needles and vials. There may be special provincial or local laws for disposal of used needles and syringes.

- Place all used needles, needle covers, syringes, and vials (empty or unused contents) into a “Sharps” container given to you by your doctor or pharmacist or in a hard-plastic container with a screw-on cap, or a metal container with a plastic lid, labelled “used syringes.” Do not use glass or clear plastic containers.
- When the container is full, tape around the cap or lid to make sure the cap or lid does not come off. **Do not throw the container in the household trash. Do not recycle.**
- **Always** keep the container out of the reach of children.

### **Usual dose:**

The usual dose of XGEVA is 120 mg administered once every 4 weeks. If you are being treated for giant cell tumour of bone or hypercalcemia of malignancy, you will receive an additional dose 1 week and 2 weeks after the first dose in the first month of treatment only.

You should also take supplements of calcium and vitamin D as instructed by your doctor.

### **Overdose:**

If you think you, or a person you are caring for, have taken too much XGEVA, contact a healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

### **Missed Dose:**

If you miss a dose you should try to receive that dose as soon as you can. In order for XGEVA to work properly, XGEVA needs to be given every 4 weeks. Continue to schedule your doses every four weeks.

### **What are possible side effects from using XGEVA?**

These are not all the possible side effects you may have when taking XGEVA. If you experience any side effects not listed here, tell your healthcare professional.

- Low blood calcium (hypocalcemia)  
Symptoms of low blood calcium may include muscle spasms, twitches, cramps, numbness or tingling in fingers, toes or around the mouth
- Skin infection with swollen, red area of skin that feels hot and tender and may be accompanied by fever (cellulitis)
- Sore in mouth involving gums or jaw bones (osteonecrosis of the jaw)
- Shortness of breath (dyspnea)
- Low phosphate levels in the blood (hypophosphatemia)
- Allergic reactions (eg, rash, hives, or in rare cases, swelling of the face, lips, tongue, throat, or trouble breathing)
- Unusual thigh bone fractures (atypical femoral fracture)
- Rash that may occur on the skin or sores in the mouth (lichenoid drug eruption)
- Hair loss (alopecia)
- Pain, sometimes severe, in the muscles, joints, arms, legs or back
- High calcium levels in the blood (hypercalcemia) after stopping treatment in patients who are still growing while on treatment with XGEVA
- High calcium levels in the blood (hypercalcemia) after stopping treatment in patients with giant cell tumour of the bone
- Broken bones in your spine (multiple vertebral fractures) after stopping treatment with XGEVA

Serious side effects and what to do about them			
Symptom / effect <sup>a</sup>	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<b>VERY COMMON</b>			
Low calcium levels in the blood [muscle spasms, twitches, cramps, numbness or tingling in fingers, toes or around the mouth] (hypocalcemia)		√	
<b>COMMON</b>			
Sore in mouth involving gums or jaw bones (osteonecrosis of the jaw)		√	
<b>UNCOMMON</b>			
Skin infection (mainly cellulitis) leading to hospitalization		√	
High calcium levels in the blood (hypercalcemia) after stopping treatment in patients with giant cell tumour of bone		√	
Unusual thigh bone fractures (atypical femoral fracture)		√	
<b>RARE</b>			
Allergic reactions [e.g. rash, hives, or in rare cases, swelling of the face, lips, tongue, throat, or trouble breathing] (drug hypersensitivity)		√	
High calcium levels in the blood (hypercalcemia) after stopping treatment in patients who are still growing while on treatment with XGEVA		√	
Broken bones in your spine following discontinuation of XGEVA (multiple vertebral fracture)		√	

<sup>a</sup> Frequency reflects all adverse events

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

### **Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

### **Storage:**

Keep out of the reach and sight of children.

XGEVA should be stored in the refrigerator at 2°C to 8°C in the original carton to protect it from light. Do not freeze XGEVA. Do not shake XGEVA.

XGEVA may be removed from the refrigerator and allowed to reach room temperature (up to 25°C) to make the injection more comfortable.

Once removed from the refrigerator, XGEVA must not reach temperatures above 25°C and must be used within 30 days. If not used within 30 days, XGEVA should be discarded.

Do not use XGEVA after the expiry date which is printed on the carton and label. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines that are no longer required.

### **If you want more information about XGEVA:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website [[www.xgeva.ca](http://www.xgeva.ca)], or
- Call Amgen Canada Medical Information at 1-866-502-6436, or
- Call the Victory Program at 1-888-706-4717.

This leaflet was prepared by Amgen Canada Inc.

Last Revised: Dec 21, 2020